



HAL
open science

Novel approach for testing the food limitation hypothesis in estuarine and coastal fish nurseries

Adrien Tableau, Hervé Le Bris, Erwan Saulnier, Olivier Le Pape, Annick Brind'Amour

► To cite this version:

Adrien Tableau, Hervé Le Bris, Erwan Saulnier, Olivier Le Pape, Annick Brind'Amour. Novel approach for testing the food limitation hypothesis in estuarine and coastal fish nurseries. *Marine Ecology Progress Series*, 2019, 629, pp.117-131. 10.3354/meps13090 . hal-02611720

HAL Id: hal-02611720

<https://institut-agro-rennes-angers.hal.science/hal-02611720>

Submitted on 14 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Novel approach for testing the food limitation hypothesis in estuarine and coastal fish nurseries

Tableau Adrien ^{1,2,3,*}, Le Bris H ¹, Saulnier E ^{1,2}, Le Pape Olivier ¹, Brind'Amour Anik ²

¹ ESE, Ecology and Ecosystem Health, Agrocampus Ouest, INRA, 65 rue de Saint-Brieuc CS 84215 35 042 Rennes Cedex, France

² Ifremer, EMH, rue de l'île d'Yeu, BP 21105, 44311 Nantes Cedex 03, France

³ University of Rhode Island, Graduate School of Oceanography, South Ferry Road, Narragansett, RI 02882, USA

* Corresponding author : Adrien Tableau, email address : adrien.tableau@gmail.com

Abstract :

Survival success of early life stages of fish is regulated by density-dependent effects, but the limiting factors explaining these effects have not been well identified. In coastal habitats, juveniles of many fish species occur in high concentrations and possibly compete for food resources. This study compared the ratio of food consumption to benthic prey production to test whether food availability is a major factor defining the carrying capacity of fish nurseries. We developed a method to quantify the exploitation efficiency (also called ecotrophic efficiency) of the juvenile fish feeding on benthic prey, expressed as a ratio of food consumption to food production. This method includes many sources of uncertainty and a key parameter of prey accessibility. Applied to the case study of the Bay of Vilaine (north Bay of Biscay), results suggest food limitation for juvenile fish. The generic nature of this method supports its wide application in various nursery habitats. As food limitation is a density-dependent process having a dampening effect on recruitment variability, examining its occurrence over time and space will improve our comprehension of nursery-dependent fish dynamics.

Keywords : Exploitation efficiency, Predator-prey relationship, Food consumption, Fish juvenile, Nursery habitat, Bioenergetics, Density dependence, Carrying capacity, Secondary production

1. INTRODUCTION

Estuarine and coastal habitats are suitable nurseries for a large proportion of demersal and benthic fish species (Seitz et al. 2014, Brown et al. 2018) and are thus considered essential for completing their life cycle (Peterson et al. 2000, Beck et al. 2001, Fulford et al. 2011). Coastal nursery dependent species congregate at the juvenile stage in these spatially restricted habitats that supply abundant food resources and act as a potential refuge from predators (Gibson 1994, Gibson et al. 2002). This fish concentration is suspected to result in density-dependent processes, limiting the growth (Bacheler et al. 2012), survival, and thus recruitment of fish populations (Iles & Beverton 2000, Minto et al. 2008). The positive relationship between surface areas of nursery habitats and the resulting average recruitment (Rijnsdorp et al. 1992, Kostecki et al. 2010, Wetz et al. 2011) supports this hypothesis. However, demonstration of density dependence does not indicate the cause of the limitation (Sinclair & Pech 1996). Even though some critical environment conditions can severely impact the carrying capacity of nursery habitats (e.g. van der Veer et al. 2000), the underlying processes remain poorly understood. Competition for food is assumed to be a key process (i.e. Gibson 1994), but the causal link between density-dependent survival and food is still debated (Le Pape & Bonhommeau 2015). Some studies suggest that benthic invertebrate populations are only minimally impacted by their predators (e.g. Gee et al. 1985, Shaw & Jenkins 1992); consequently, survival of juvenile fish would not be dependent on food availability (Ross 2003, Hampel et al. 2005, Vinagre & Cabral 2008). However, other studies propose that the distribution of benthic invertebrates can be regulated by their predators in estuarine and coastal ecosystems. Food availability would then be a factor limiting the survival of juvenile fish in nursery habitats (Nash & Geffen 2000, Nash et al. 2007, Craig et al. 2007): this is the food limitation hypothesis (Le Pape & Bonhommeau 2015).

Two approaches are commonly used to test whether food is limiting in coastal areas. The indirect approach focuses on growth variations. It assumes that the *in situ* growth rate would decrease with the density of juveniles if they compete for food (e.g. van der Veer & Witte 1993, Craig et al. 2007). However, this approach could lead to the erroneous rejection of the hypothesis of food limitation. Indeed, even in a food limitation context, the apparent growth of a juvenile fish population could remain maximal because of the highly size-selective mortality of juvenile fish: larger individuals have a higher survival than smaller ones (Sogard 1997, Levin et al. 1997, Le Pape & Bonhommeau 2015). Alternatively, the direct approach quantifies the exploitation pressure applied by consumers on their prey. Exploitation efficiency (EE; also called ecotrophic efficiency) is the ratio of food consumed by the juvenile fish community to the food supply represented by the production of the benthic invertebrate community (Collie 1987, Vinagre & Cabral 2008). This ratio is an intermediate component to compute ecological efficiency, which quantifies energy transfer among trophic levels (Lindeman 1942). It directly quantifies the portion of benthic production consumed by their predators; thus, it is an indicator of predation pressure. In this context, the food limitation hypothesis would be accepted if EE is high over years and/or over nursery habitats.

Few studies have carried out the direct approach to assess the predation pressure applied by one ecological guild on another. This method requires collecting a large amount of quantitative data on many marine species, including both the marine juvenile fish community and their benthic prey. Generally, one of the 2 following methods are used to estimate food consumption (FC). First, the gut-content method, which is a flux analysis requiring time-consuming experimental work. Gut contents must be weighed at constant intervals to determine the evacuation rate of each fish species (Elliott & Persson 1978). This method likely underestimates FC, as regurgitation often occurs soon after fish capture (e.g. Elliott & Hemingway 2002, Vignon & Dierking 2011). The second approach is based on bioenergetics and estimates FC from fish growth. The energy conversion process can be more or less exhaustive; e.g. ranging from a very detailed approach using bioenergetics models (e.g. dynamic energy budget [DEB] theory; Kooijman 2009) to a largely simplified one, using mass-balanced trophic models (e.g. the Ecopath framework; Christensen et al. 2005). Assessing the quantity of food that is ingested to obtain annual fish production requires a compromise between (1) an exhaustive description of the processes involved in the conversion of ingested food into fish biomass and (2) the availability of reliable data to describe this conversion process. This compromise precludes the systematic use of bioenergetics models. DEB models have been used successfully to estimate juvenile food limitation for a well-studied coastal nursery dependent species (van der Veer et al. 2010, 2016). However, the DEB approach requires data and knowledge on a variety of fine metabolic processes that are mostly undescribed

for marine juvenile fish, preventing its use in a multispecific approach. Mass-balanced trophic models are used to estimate flows between functional groups in an ecosystem (Christensen et al. 2005). These models, also based on bioenergetics, estimate the total EE of prey. However, mass-balanced models use an oversimplified description of growth and are not well suited to analyze short-term processes (i.e. turnover rate of prey production in coastal and estuarine systems; Ritter et al. 2005). Hence, these 2 families of models are not designed to focus on the food limitation hypothesis in estuarine and coastal fish nurseries. Therefore, we need a method that is based on bioenergetics, but less complex than a full DEB calculation.

The few studies that focus on predation of benthivorous fish generally estimate a low EE of the benthic prey, indexing low predation pressure (e.g. Vinagre & Cabral 2008). However, they do not necessarily conclude that there is an absence of food limitation (Collie 1987). Indeed, benthic production is shared by many fish species in coastal nursery habitats (McLusky & Elliott 2004, Nicolas et al. 2010). By considering only a few consumer species, most studies assess only a small part of the total EE of prey production. This key point underlines the need to account for a representative part of the consumers relying on the same pool of prey in order to fully evaluate the food limitation hypothesis.

The limits of the existing approaches to assess fish consumption emphasize the need to develop a bioenergetics-based approach responding to the constraints induced by data availability, but also sufficiently precise to quantify predation pressure and thus deal with the food limitation hypothesis in multispecies fish nurseries. In this paper, we developed a method to quantify energy flow between 2 compartments: the benthic invertebrate community and the benthivorous juvenile fish community. This methodology aims to estimate the ratio of FC of juvenile fish to benthic food production (FP) by quantifying 2 key intermediate components: fish production and prey production (Fig. 1). This approach accounts for prey accessibility and quantifies the uncertainty associated with input parameters. We applied this methodology to the Bay of Vilaine nursery ground (Bay of Biscay, Western Europe) to test the food limitation hypothesis.

2. MATERIALS AND METHODS

2.1. Computing EE

Our objective was to quantify energy flow between benthic macroinvertebrates and benthivorous fish living in estuarine and coastal nursery ecosystems, a community mostly dominated by juvenile fish (Gibson 1994). We estimated EE, which is the ratio of juvenile fish FC to benthic FP. EE is a baseline for the analysis of the food limitation hypothesis in fish nursery habitats (Fig. 1).




Fig. 1. Computational flowchart used to estimate exploitation efficiency (EE), including uncertainty analysis with Monte Carlo sampling. AFP: accessible food production

2.2. FC

The FC metric estimates the quantity of food consumed annually by a community of consumers. The bioenergetics approach computes fish production to then determine the FC of juvenile fish. For a single individual, annual production refers to its growth over 1 yr. Extrapolation to the population level is carried out either by integrating over time the product between instantaneous fish number and instantaneous growth rate, or by using its production-to-biomass ratio, which integrates both. Unlike macrobenthic invertebrates, a large proportion of fish species are not resident in coastal shallow water habitats, as they migrate to deeper waters as adults (Seitz et al. 2014). In this context, the production-to-biomass ratio, referring mostly to an entire population in the literature, is not appropriate to estimate juvenile fish production within nursery habitats. Moreover, as fish feed on benthic macroinvertebrates for only a limited period of the juvenile stage, the time window must be properly delimited. Both arguments lead to estimate FC from mortality and growth rates. Data collection is detailed in Supplement S2 at www.int-res.com/articles/suppl/m629_p117_supp.pdf.

The biomass production (P , in weight yr^{-1} or $\text{weight surface}^{-1} \text{yr}^{-1}$) of a juvenile cohort during the growth period is estimated as:

$$P = \int_{t=t_0}^{t=T} n(t) \cdot \frac{dw}{dt} \cdot dt \quad (1)$$

where t_0 and T , respectively, refer to the beginning and end of the growth period when juvenile fish feed on benthic macroinvertebrates and water temperature is correspondingly higher; $n(t)$ is the number of individuals at time t ; and $\frac{dw}{dt}$ is the instantaneous growth rate. As the integration in Eq. (1) is not straightforward, it is discretized through a daily basis in Eq. (2) to estimate P of a juvenile cohort (in weight yr^{-1} , or in $\text{weight surface}^{-1} \text{yr}^{-1}$):

$$P = \sum_{d \in d_0:D} n_d \cdot (w_d - w_{d-1}) \quad (2)$$

where d_0 and D , respectively, are the first and last day of the growth period; n_d is the number of individuals in the middle of day d ; and w_d and w_{d-1} are the individual weights at the end of day d and day $d-1$. The difference

between the 2 weights provides the individual production for day d. The 2 components of Eq. (2) (n_d and w_d) can be computed using survey data, daily growth rate, and daily mortality rate. n_d is estimated as:

$$n_d = \frac{c_s}{q} \cdot e^{-(d-s)Z} \quad (2.1)$$

where s is an index of the survey date; c_s is fish abundance in number or number surface⁻¹; q is catch efficiency; and z is daily mortality (in d⁻¹). w_d is estimated as:

$$w_d = a \cdot [L_s + (d + 1/2 - s) \cdot G]^b \quad (2.2)$$

where G is the daily growth rate of the studied species observed in the literature (Supplement S2); L_s is the average fish length of a single cohort observed in the survey at day s; and a and b are parameters of the length-weight relationship. d_0 (Eq. 2.3) is the day when an individual fish reaches the minimum size to actually feed on macrofauna:

$$d_0 = \frac{L_{d_0} - L_s}{G} + s \quad (2.3)$$

where G (defined above) is assumed to be constant during the main growth period; L_s is as defined above; and L_{d_0} is the average fish length corresponding to a diet shift towards macroinvertebrates, defined from gut content analyses (Tableau et al. 2015). P can then be expressed from Eqs. (2), (2.1), (2.2), and (2.3) as follows:

$$P = \frac{c_s}{q} \cdot e^{sZ} \cdot a \cdot \sum_{d \in d_0:D} e^{-dZ} [(L_s + (d + 1/2 - s) \cdot G)^b - (L_s + (d - 1/2 - s) \cdot G)^b] \quad (3)$$

where D, the end of the main growth period, occurs in late fall in temperate ecosystems (Hamerlynck & Hostens 1993). FC is then derived from P. As the bioenergetics approach refers to energy units, P must be converted into energy production. To do so, we used data on gross conversion efficiency (K) (Hidalgo et al. 1987), defined as the quantity of ingested energy required to produce one energy unit of juvenile fish. As macroinvertebrates may represent only a part of the diet of some roundfish species (Hamerlynck & Hostens 1993), only that part of the production must be considered. Therefore FC of the benthivorous fish community, expressed in energy yr⁻¹ or energy surface⁻¹ yr⁻¹, is calculated as:

$$FC = \sum_{i \in 1:1^{th} \text{ fish cohort}} P_i \cdot DC_i \cdot E_i \cdot \frac{1}{K_i} \quad (4)$$

where P_i is the biomass production of fish cohort i; DC_i is the proportion of benthic macroinvertebrates in the diet estimated using stomach contents; and E_i is the energy density gathered from the literature (Supplement S2).

2.3. Benthic FP

Benthic invertebrate species were considered potential prey if they have been observed in juvenile fish gut contents (determined from literature studies on juvenile diet or from our own observations; see details in Tableau et al. 2015 and Supplement S2). Since benthivorous juvenile fish are considered to be opportunistic feeders (De Vlas 1979, Hampel et al. 2005, van der Veer et al. 2011, Schückerl et al. 2012), FC cannot be partitioned among prey species. For consistency, the biological productions of all macrobenthic prey species are summed to estimate FP. As the mobility of benthic invertebrate prey is limited, one can assume that they remain in the coastal nursery all their lives. Thus, production-to-biomass ratio data referring to entire populations can be used to compute their production (Brey 2001). The 'available benthic energy coefficient' (ABEC) was developed for this purpose (Tableau et al. 2015). The Brey Model was used because it the most accurate model, especially for marine assemblages (Brey 2012). Since the benefits of a correction coefficient suggested in Brey (2012) remain an open question, we chose not to use it. As ABEC is only applicable to average annual biomasses, biomasses observed during the survey were corrected. The resulting biomass production was converted into energy by a species-specific coefficient, since energy density varies widely among in vertebrate species. For instance, energy density is 1.02 kJ g⁻¹ for the bivalve *Corbula gibba*, whereas it is 5.84 kJ g⁻¹ for the polychaete *Lumbrineris* sp. and 8.19 kJ g⁻¹ for *Ampelisca* sp. (Brey et al. 2010). FP is expressed in energy yr⁻¹ or energy surface⁻¹ yr⁻¹ as follows:

$$FP = \sum_{j \in 1:j^{\text{th}} \text{ prey species}} CR \cdot B_j \cdot P \cdot B_j \cdot (1 + R_j) \cdot E_j = \sum_{j \in 1:j^{\text{th}} \text{ prey species}} CR \cdot B_j \cdot ABEC_j \quad (5)$$

where CR is a ratio converting the biomass observed during a survey at a given period of the year into average annual biomass (Tableau et al. 2015); B_j is the biomass observed during a survey (in weight or weight surface⁻¹); $P \cdot B_j$ the production-to-biomass ratio estimated from an empirical model accounting for water temperature, but which does not account for regeneration after sublethal fish cropping (Brey 2012) (in yr⁻¹); R_j is a regeneration coefficient (Tableau et al. 2015) accounting for somatic regeneration (i.e. production-to-biomass must be raised by 15% for *Amphiura filiformis* to account for regeneration; Skold et al. 1994); and E_j is the energy density (in energy weight⁻¹) for the j^{th} prey species.

Only accessible prey matters for a predator (Tableau et al. 2015, van der Veer et al. 2016). ABEC (Tableau et al. 2015) allows us to weight the FP of a species by an accessibility coefficient based on observed juvenile fish prey catch rates. According to Tableau et al. (2015), 2 accessibility categories can be statistically identified; the authors suggested setting 1 as the coefficient of easily accessible prey and 0.11 as the coefficient of hardly accessible prey. When included in the FP equation (Eq. 5), the accessibility coefficient (A_j) gives the accessible FP (AFP):

$$AFP = \sum_{j \in 1:j^{\text{th}} \text{ prey species}} CR \cdot B_j \cdot P \cdot B_j \cdot (1 + R_j) \cdot E_j \cdot A_j \quad (6)$$

2.4. Partial uncertainty analyses

The computation of EE requires a large amount of data and parameters, usually at the species level (Table 1). Since quality of information can be poor on some parameters and data, uncertainty analyses were carried out. To do so, we classified the data and parameters following their quality in 3 categories, each of which were treated differently (Table 1).

- 1st category: data and parameters with negligible uncertainty. Information comes from the studied species. Values are defined as fixed values. Typically, fish length can be considered in this category.
- 2nd category: data and parameters with highly quantified uncertainty. Information comes from the studied species but is uncertain given observation error (e.g. data on density), given the region and/or life stage at which the information is gathered, or because it is inferred from species taxonomically and functionally close to those observed in the data set (e.g. parameters on mortality rate). For this category, uncertainty is considered as known and/or estimated from the different sources of information. Each data point or parameter is defined using a statistical distribution (see Table 1 and Supplement S1 for details on the distributions).
- 3rd category: data and parameters with highly unquantified uncertainty. Unlike the 2nd category, there is a lack of knowledge and references to inform the values of this category. A conservative approach is needed. It consists of selecting, in a large range of uncertainty, the value of the data or parameter leading to a minimal EE. When at least one parameter is defined in that category, only a minimum threshold estimate of the EE is estimated. EE was computed using the Monte Carlo method (Metropolis & Ulam 1949). Random sampling in parameter distributions were repeated 5000 times to build the final distributions (Fig. 1).

Table 1. Categorization for the data and parameters. A resolution at the species level (yes, Y) associated with negligible uncertainty (Y) leads to the use of fixed values. When data and parameters are not defined at the species level (no, N) and/or have high uncertainty (N), there are 2 possibilities: if several sources are available (Y), values are defined with an associated distribution; if the uncertainty is unknown (N), a conservative approach is used. N*: data are unavailable for some fish cohorts, but the latter are still informed using data from other fish species. /: not applicable

Parameter	Description	Sources of information	Information at species level	Negligible uncertainty	Known uncertainty	Category of uncertainty	Statistical distribution
Food consumption in energy (fish data)							
C _s	Fish abundance	} Surveys	Y	N	Y	2-distribution	Gamma
s	Survey date index		Y	Y	/	1-fixed data	/
L _s	Fish length	} Deduced from surveys	Y	Y	/	1-fixed data	/
a	Length-weight parameter		Y	Y	/	1-fixed data	/
b	Length-weight parameter		Y	Y	/	1-fixed data	/
L _{d0}	Diet shift length		Y	Y	/	1-fixed data	/
d ₀	Diet shift date index	} Deduced from surveys and literature	N*	N	N	3-conservative	/
D	Growth end index		N	N	N	3-conservative	/
q	Catch efficiency	} Literature	N*	N	N	3-conservative	/
Z	Mortality rate		N*	N	N	3-conservative	/
G	Growth rate		N*	N	N	3-conservative	/
DC	Diet composition		Y	N	N	3-conservative	/
E	Energy density		N	N	Y	2-distribution	Gamma
K	Gross efficiency		N	N	Y	2-distribution	Inverse gamma
Food production in energy (prey data)							
B	Invertebrate biomass	Surveys	Y	N	Y	2-distribution	Gamma
CR	Conversion ratio	} Deduced from surveys	N	N	Y	2-distribution	Beta
A	Accessibility		N	N	Y	2-distribution	Beta
P:B	Production-to-biomass ratio	} Literature	Y	N	Y	2-distribution	Lognormal
R	Regeneration rate		N	N	N	3-conservative	/
E	Energy density		Y	Y	/	1-fixed data	/

2.5. Application to a temperate nursery habitat: the Bay of Vilaine, France

2.5.1. Study site

The Bay of Vilaine is a soft-bottom habitat under estuarine influence that is used as a nursery by several benthic and demersal fish species of commercial interest (Le Pape et al. 2003). It has been studied for more than 30 yr, with valuable knowledge gained regarding its fish (Nicolas et al. 2007, Kopp et al. 2013) and benthic invertebrate communities (Le Bris & Glemarec 1995, Brind'Amour et al. 2014). Given such data and knowledge-rich context, the bay constitutes a relevant framework in which to analyze predator-prey relationships (Le Pape et al. 2003, Kostecki et al. 2010, Tableau et al. 2016). The studied area covers the subtidal zone located from 5 to 30 m depth and consists mostly of sandy mud sediments (Le Bris & Glemarec 1995). It was stratified into 3 zones along the bathymetric gradient (see Supplement S2 for details). All samples were collected exclusively on this type of sediment, explaining the shape of the geographic domain of the study (Fig. 2).




Fig. 2. Sampling design in the Bay of Vilaine. Red dots and blue crosses: grab and trawl sampling stations, respectively; 3 zones: the 3 main subtidal soft-bottom habitats along the bathymetric gradient. The C-shape of the area is due to a non-sampled rocky habitat on the east side of the bay

2.5.2. Survey data

A survey was performed in late September 2008 targeting juvenile fish and benthic macroinvertebrates. This season was selected because, for most of the local fish species, juveniles born in spring are large enough to avoid selectivity issues and size-dependent catch efficiency, and also because this period is just before their autumnal migration towards deeper grounds after the growth period (Dorel et al. 1991, Le Pape et al. 2007). Both protocols are detailed in Tableau et al. (2015). For fish sampling, most of the catches corresponded to benthic and demersal juvenile fish species belonging to the age groups 0 and 1 yr (G0 and G1). Fish species feeding at least partially on benthic macroinvertebrates were selected for this case study: 3 flatfish (benthic) species (*Solea solea*, *Dicologlossa cuneata*, and *Pleuronectes platessa*) and 4 roundfish (benthic or benthopelagic) species (*Merlangius merlangus*, *Merluccius merluccius*, *Mullus surmuletus*, and *Trisopterus luscus*). They represent 71% of the total fish abundance in this area. Ninety-four taxa were considered potential prey for the benthivorous fish community based on gut content analyses. They represent 96% in weight of the collected benthos. See Tableau et al. (2015) for further details on that selection.

2.5.3. Data inputs

Data and parameters are shown in Table 1. The detailed methodology for each data component is fully detailed in Supplement S2. Out of 14 parameters used to compute FC, 6 are described using the conservative approach, i.e. we select the value of the data or parameter leading to a minimal EE. For FP, 1 parameter out of 6 used the conservative approach.

3. RESULTS

3.1. FP and FC

The total FP in 2008 averaged over the 3 zones was 8120 MJ ha⁻¹ yr⁻¹ (~812 kJ m⁻² yr⁻¹) and varied from 5850 to 8620 MJ ha⁻¹ yr⁻¹ among zones (Fig. 3a). When that FP is weighted by the accessibility coefficient (i.e. AFP), it is lowered to an average of 1110 MJ ha⁻¹ over the 3 zones (Fig. 3b), corresponding to 14% of total FP.

The spatial distribution of the FC (Fig. 3c) is more contrasted than the AFP (e.g. 2 times higher in zone 3 than in zone 2). Total FC in 2008 averaged among the 3 zones is estimated at 485 MJ ha⁻¹ yr⁻¹ for the whole bay.

The main consumers are 2 exclusive benthivorous fish cohorts (G1 [1-yr age group] of *Solea solea* and G1 of *Dicologlossa cuneata*) and 2 partial benthivorous cohorts (G0 of *Trisopterus luscus*, and G0 of *Merlangius merlangus*). The consumption of the G0 of *S. solea* and G0 of *Pleuronectes platessa* is concentrated in zone 1, which is a shallow area (5 to 13 m depth).




Fig. 3. (a) Total and (b) available food production and (c) food consumption, displayed by fish cohort (G0: young of the year; G1: 1 yr old) in the bay of Vilaine. Boxes: 50% confidence intervals; lines: 95% confidence intervals

3.2. EEs

The EE (Fig. 4) of the main benthivorous species (*S. solea*) on total FP is 2.1%. When all benthivorous species are considered, the estimate is 3.5% and increases to 6.1% when partial benthivorous species are also included. When only AFP is considered, the estimate of EE is far higher, ranging from 17.7% for *S. solea* to 50.0% for all species. Uncertainties around these estimates are very large, especially when only AFP is considered.




Fig. 4. Exploitation efficiency of total and accessible benthic production in 2008. Exclusive benthivorous species are the flatfish *Solea solea*, *Dicologlossa cuneata*, and *Pleuronectes platessa* and the roundfish *Mullus surmuletus*. All benthivorous species include also *Merluccius merluccius*, *Merlangius merlangus*, and *Trisopterus luscus*. Boxes: 50% confidence intervals; lines: 95% confidence intervals

4. DISCUSSION

This study demonstrates a method of computing the EE of fish juveniles feeding on benthic prey in nursery habitats. Properly estimating EE is essential to better understand the functioning of marine ecosystems (Libralato et al. 2008, Watson et al. 2014), and more specifically, to investigate the food limitation hypothesis within fish nursery habitats (Le Pape & Bonhommeau 2015).

4.1. The food limitation hypothesis: preliminary insights

Coastal nursery habitats present generally high FP and a refuge from predators (Wouters & Cabral 2009, Ryer et al. 2010, Nash & Geffen 2012). The underlying aim of the food limitation hypothesis is to determine predation pressure in order to understand the level of food limitation (Le Pape & Bonhommeau 2015). Defining different food limitation scenarios can help in interpreting the EE value for juvenile fish:

- No food limitation: food is abundant, and juvenile fish can invest negligible time and energy in foraging. The size of the juvenile population is either defined by factors affecting mortality in previous life stages (e.g. spawner condition affects offspring condition and survival, Hare 2014; larval predation, Houde 2008) or by other factors that occur during the juvenile stage (e.g. predation mortality, competition for space, Sheaves et al. 2015). Under the 'no food limitation' condition, the EE (EE_0) varies over time (inter-annual independence between FP and FC) but remains close to 0. The reciprocal to this causal relationship is not always true: EE can be close to 0 at the scale of the entire nursery with food limitation that occurs locally if there is a lot of spatial heterogeneity of fish and prey distributions.
- 1st level of food limitation: FP supports normal growth of the juvenile fish community, but overdispersed prey force the fish to invest a significant amount of time and energy in foraging. This behavioral response increases the vulnerability of juvenile fish and, consequently, their mortality (Gibson 1994, Biro et al. 2003). This EE (EE_1) is significantly above 0, but depends on density-dependent predation pressure (Johnson 2007, Cebrian 2015). This limiting effect, called predation-sensitive foraging (Sinclair & Arcese 1995) or risk effect (Heithaus et al. 2008), is typically a combination of food limitation and predation.
- 2nd level of food limitation: FP is temporarily not sufficient for all fish juveniles (van der Veer et al. 2016). Juvenile fish must invest a lot of time in finding food, and thus instantaneous vulnerability is exacerbated. One main difference between this and the 1st level of food limitation is that some individuals have a lower body condition and remain small. Paradoxically, a decrease of the apparent growth of the population is rarely observed even though this level of food limitation is suspected to occur frequently (Walters & Juanes 1993, Fiksen & Jørgensen 2011, Le Pape & Bonhommeau 2015). Indeed, small size implies vulnerability to predators that leads to size-dependent mortality (Sogard 1997), the apparent growth is thus maintained at a maximum. If this level of food limitation unfaillingly occurs, the EE (EE_2) consistently, and over years, reaches a higher value than EE_1 (which is, however, still unknown).
- 3rd level of food limitation: long-term FP is insufficient for juvenile fish. Consequently, individuals frequently starve to death (Gibson 1994). This has been verified in experiments (Edwards et al. 1970), but has not been observed in natural environments. Indeed, food deprivation in the wild, leading to poor condition, increases vulnerability to diseases and predation (Vethaak 1992) and leads to rapid and massive mortality (Juanes 2007) in the short term. This EE (EE_3) is expected to reach values near 100% if all benthic invertebrate consumers are considered or if the EE is estimated from AFP when only juvenile fish are considered.

Consequently, juvenile fish communities are affected by the lack of food availability before full exploitation at the scale of the fish nursery (Heath 2005). Moreover, low EE does not exclude a high level of food limitation as predicted by the foraging arena theory (Walters & Juanes 1993). Indeed, high predation pressure on fish juveniles prevents them from leaving their shelter to reach better feeding areas and forces them to feed in very restricted areas that are then potentially overexploited. Nevertheless, as nursery habitats are considered to be shelters for juvenile fish (Gibson 1994, Wouters & Cabral 2009), EE could potentially be high. Thus, the remaining problem is to assess the thresholds of EE above which food limitation is temporarily (2nd level) or ongoingly (3rd) insufficient (i.e. EE_2 and EE_3 , respectively). The 3rd level could be reached before an EE of 100%. Indeed, significant levels of exploitation below 100% could lead to a collapse in prey availability. For instance, the uptake of 50% of the available production in an ecosystem (e.g. combination of invertebrate species and shellfish farming on primary production; Arbach Leloup et al. 2008) could lead to a dramatic shift in the trophic chain and a strong limitation of food availability for natural consumers (Raimonet & Cloern 2017).

Moreover, considering the widely studied fisheries exploitation, the sustainable level of EE at an ecosystem scale is estimated to be around only 15% (Libralato et al. 2008, Watson et al. 2014, Zhou et al. 2015). At this rate, transfer efficiency of benthic production in shallow coastal and estuarine nursery grounds could be limited (Arbach Leloup et al. 2008, Raimonet & Cloern 2017). The specificities of the communities of these ecosystems, i.e. the quality paradox (Elliott & Quintino 2007), with a large number of immature individuals and high levels of mortality, leads to moderate sustainable levels of transfer losses, and thus low potential EE. As these considerations concern overall EE, including predation by competitors, the maximal level of EE by juvenile fish is probably dramatically lower (see Table 2), far below 100%.

The estimates established in the Bay of Vilaine suggest that at least the 1st level of food limitation occurred in 2008. The spatial overlap between the biomass of the juvenile fish community and FP (macrobenthic prey) over the Bay of Vilaine (Tableau et al. 2016) supports this hypothesis. The high EE estimate on AFP suggests that the 2nd level of food limitation might be reached in some sectors of the bay. The potentially large underestimation of the level of EE and previous considerations on the likely moderate threshold of lasting food limitation (i.e. EE₃) lead us to conclude that the 3rd level of food limitation could potentially be reached. An investigation over several years would be required to determine if there is consistency over time. The large uncertainties estimated in the present study would be drastically reduced by conducting at least 2 surveys (at the beginning and end of the main growth period) instead of one, as this increase in data would provide better estimates of fish growth, fish mortality (Nash & Geffen 2000), average benthic invertebrate biomass (Saulnier et al. 2019), and accessibility coefficients. Even if the food limitation hypothesis is investigated at the fish community scale, we assume that the density- dependent response to a lack of food availability would vary from one fish species to another. For instance, we expect that an exclusive benthic feeder would be more sensitive to a lack of benthic invertebrate availability than a partial benthic feeder, which can adapt its diet.

4.2. Conservative (under)estimation of EE

The proposed method provides an estimate of the EE that integrates available ecological knowledge about the various species included in this case study. Input values can be derived either from *in situ* data (whenever possible) or from the literature. With regards to the assessment of their reliability, they are either fixed values or defined as statistical distributions. Unfortunately, available information on some uncertain parameters is too scarce and/or unreliable to define a statistical distribution. This limitation was overcome by using a conservative approach. Parameter values were chosen to provide a minimum estimate of FC. Improving these parameters would bring the estimate closer to the actual EE. Several other shortcomings of the approach lead to an underestimation of EE:

- Fish biomass: the use of a beam trawl minimizes the uncertainty of benthic and juvenile fish catch per trawled area (Dorel et al. 1991). However, estimation methods used in the literature often only provide maximum catch efficiencies (e.g. Reiss et al. 2006), thus minimum estimates of fish biomass. Moreover, fish species selected in the analysis account for 71% of the catch, not the overall amount that fish consumed.
- One parameter using the conservative approach is diet composition. Macrobenthic proportion in the diet is defined with the smallest values observed in the literature (Supplement S2). A source of underestimation of FC is that some fish cohorts are present in the bay only during one season. Consequently, they are not observed in some surveys, and are therefore not considered (e.g. G1 of *Pleuronectes platessa* in early summer, Desaunay et al. 1981). Moreover, the considered growth period is restricted to a time window corresponding to only the maximum growth period (Hamerlynck & Hostens 1993). Feeding activity during the winter period is not estimated. Juveniles of the selected species move progressively to deeper water in winter, but the corresponding consumption during this migration cannot be estimated as the apparent mortality rate (reflecting both mortality and migration rates) is not available. Consumption by the resident fish species, present in the remaining 29% of the survey catch, likely occurs (Fonds et al. 1992).
- The consumption of benthic production was estimated for fish species only, whereas other non-fish consumers (e.g. the brown shrimp *Crangon crangon*), sometimes occurring at high biomass, may also feed on the same pool of prey (Jung et al. 2017).
- The parameter converting energy production into energy consumption is defined and does not account for indigestible parts of ingested prey.

All of these shortcomings most likely lead to an underestimate of FC occurring in the natural environment. The actual EE value is higher than the estimate provided. This (under)estimate is valuable in the context of the food limitation hypothesis, but associated conclusions need to account for this conservative methodology.

4.3. Community scale and prey accessibility: 2 keys of investigation

The 2008 FP in the bay of Vilaine is in the production range of similar habitats (Burd et al. 2012, Bolam et al. 2014). When only the dominant benthivorous species is considered, the EE on the FP is low (~2%). This result is in the range of studies focusing on a single fish species (Table 2). However, since the carrying capacity of a population is determined by the strength of interspecific density-dependence (Brown et al. 2019), a community approach was adopted. When more of the benthivorous fish community was considered, the EE was far higher (~6%). This result underlines the need to account for a maximum of species preying on the same community (Vinagre & Cabral 2008). The EE of the juvenile fish of the selected species is in the range of the results of Ecopath applications in nursery habitats (Table 2). However, these models were not built to assess the food limitation hypothesis and face several shortcomings (i.e. the total biomass of a fish species is sometimes estimated from trawl catch without accounting for the catch efficiency, Rybarczyk & Elkaïm 2003); the lack of knowledge regarding production-to-biomass ratios for juvenile fish is also a major concern (Mackinson & Daskalov 2007). The present estimate of EE is also close to Pihl (1985), who used a similar approach in a more restricted area (1.5 ha; Table 2).

Pihl (1985) also stated that the production of some prey species is fully consumed and that half of total prey production is consumed (Table 2), suggesting food competition, and consequently, potential food limitation. Similarly, the Ecopath-based approach estimates the overall EE as ranging from 20 to 99%. This indicates that considering the juvenile fish community is relevant, but not sufficient, as other predators may feed on the same pool of prey. The EE estimates would thus benefit from considering additional megabenthic invertebrate consumers, such as large echinoderms (e.g. starfish *Asterias rubens*) or crustaceans (swimming crab *Liocarcinus holsatus*, and the brown shrimp *Crangon crangon*). However, some key parameters referring to these species, such as the gross conversion efficiency, are unknown, which prevents the inclusion of such consumers in this application.

Van der Veer et al. (2016, p. 105) emphasized that 'for a predator only harvestable prey matters'. Even though some prey species belong to the hyperbenthos, a large part of the benthic macrofauna live partly or totally burrowed in the sediment, so access to prey items is a factor that potentially heavily impacts the food quantity that is harvestable by juvenile fish (Tableau et al. 2015). Indeed, the EE could never reach 100% even if juvenile fish were the only consumers. Considering total FP instead of AFP would fail to test the food limitation hypothesis (Vinagre & Cabral 2008). To our knowledge, this study is the first attempt to investigate the food limitation hypothesis by accounting quantitatively for food accessibility. The application to the bay of Vilaine in 2008 shows that at least 20% of the FP accessible to juvenile fish is consumed, and does not exclude that the entire accessible production is consumed (cf. the uncertainty intervals; Fig. 3). Accounting for prey accessibility significantly increases the EE estimate by including additional information that is essential to investigate the food limitation hypothesis. The accessibility coefficient provided by Tableau et al. (2015) is, however, a large source of uncertainty and requires further investigation (e.g. seasonal prey availability and ontogenetic variations, Beukema et al. 2014, van der Veer et al. 2016; species-specific behavior, de Groot 1969).

Table 2. Estimates of exploitation efficiency (EE; ratio of benthic production consumed by their predators to the total benthic production) in temperate marine ecosystems: habitat, scale, number of species, method, habitat features (area, bathymetry) and location

Main scale production of food estimation	No. of fish cohorts / species	Method of food consumption estimation	Surface area (ha)	Bathymetry (m)	EE by fish (%)	EE by all consumers (%)	Location	Study
Nursery habitat								
Species	7	Bioenergetics	19351	5–30	6.2	?	Bay of Vilaine, France	This study
Species	3	Evacuation rate	1.5	<1	6.8	50.8	Gullrnarsvik, Sweden	Pihl (1985)
Taxonomic groups	2	Evacuation rate	2400	1.6	<1.0	?	Tagus estuary, Portugal	Vinagre & Cabral (2008)
Taxonomic groups	1	Evacuation rate	4600	4.4	<1.0	?	Tagus estuary, Portugal	Vinagre & Cabral (2008)
Ecological guilds	Whole fish community	Bioenergetics (Ecopath)	>10 000	?	24	99	Narragansett Bay, USA	Monaco & Ulanowicz (1997)
Ecological guilds	Whole fish community	Bioenergetics (Ecopath)	>10 000	?	4	99	Delaware Bay, USA	Monaco & Ulanowicz (1997)
Ecological guilds	Whole fish community	Bioenergetics (Ecopath)	>10 000	?	11	94	Cheasapeake Bay, USA	Monaco & Ulanowicz (1997)
Ecological guilds	Whole fish community	Bioenergetics (Ecopath)	650 000	0–20	10	78	Seine Bay, France	Tecchio et al. (2015)
Ecological guilds	Whole fish community	Bioenergetics (Ecopath)	450 000	0–10	1	20	Gironde estuary, France	Lobry et al. (2008)
Ecological guilds	Whole fish community	Bioenergetics (Ecopath)	5000	0 (intertidal)	6.9	76	Bay of Somme, France	Rybarczyk et al. (2003)
Ecological guilds	Whole fish community	Bioenergetics (Ecopath)	3500	0–5	18	72	Seine estuary, France	Rybarczyk & Elkaïm (2003)
Adult habitat								
Species	2	Bioenergetics	3.6	<1	9	?	Kvarnbukten Bay, Sweden	Evans (1983)

4.4. Future considerations

Moving forward, it would be fruitful to compare EE between nursery habitats over time, especially in areas with large differences in juvenile fish abundance. Indeed, as an EE value of 100% is unlikely, EEs significantly different from 0% and consistent among nursery habitats with contrasted fish juvenile abundance would support a general food limitation effect. In this context, the environmental factors influencing benthic invertebrate production may also contribute to density-dependent survival explaining recruitment variations (Salen-Picard et al. 2002). As predators of juvenile fish influence fish juvenile densities, EEs could provide indirect information about these predators. According to the foraging arena theory applied to 2 nurseries monitored over several years, the maximum EE value reached in a nursery more exposed to large predators is expected to be lower than in the more protected nursery. Therefore, the maximum EE value for a nursery over a time series would reflect the quality of shelter of that nursery from predators of the juvenile fish.

The present method was developed to test the food limitation hypothesis in a temperate coastal nursery context, but this framework is sufficiently generic and could be applied to a broad latitudinal range of coastal habitats (tropical, subarctic). However, parameters that vary with latitude (e.g. duration of growth period, Freitas et al. 2012) and/or at small spatial scales (e.g. growth rate, Ciotti et al. 2014), would have to be properly defined.

Acknowledgements. Authors thank the crew of the Research Vessel Gwen-Drez employed during the nursery scientific surveys of 2008. We also thank Jeremy Collie and 4 anonymous reviewers for their fruitful advice that improved the manuscript.

LITERATURE CITED

- Arbach Leloup F, Desroy N, Le Mao P, Pauly D, Le Pape O (2008) Interactions between a natural food web, shellfish farming and exotic species: the case of the Bay of Mont Saint Michel (France). *Estuar Coast Shelf Sci* 76: 111–120
- Bacheler NM, Buckel JA, Paramore LM (2012) Density-dependent habitat use and growth of an estuarine fish. *Can J Fish Aquat Sci* 69: 1734–1747
- Beck MW, Heck KL Jr, Able KW, Childers DL and others (2001) The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. *BioScience* 51: 633–641
- Beukema JJ, Cadée GC, Dekker R, Philippart CJM (2014) Annual and spatial variability in gains of body weight in *Macoma balthica* (L.): relationships with food supply and water temperature. *J Exp Mar Biol Ecol* 457: 105–112
- Biro PA, Post JR, Parkinson EA (2003) From individuals to populations: prey fish risk-taking mediates mortality in whole-system experiments. *Ecology* 84: 2419–2431
- Bolam SG, Coggan RC, Eggleton J, Diesing M, Stephens D (2014) Sensitivity of macrobenthic secondary production to trawling in the English sector of the greater North Sea: a biological trait approach. *J Sea Res* 85: 162–177

- Brey T (2001) Population dynamics in benthic invertebrates. A virtual handbook, version 01.2. www.thomas-brey.de/science/virtualhandbook/
- Brey T (2012) A multi-parameter artificial neural network model to estimate macrobenthic invertebrate productivity and production. *Limnol Oceanogr Methods* 10: 581–589
- Brey T, Mueller-Wiegmann C, Zittier ZMC, Hagen W (2010) Body composition in aquatic organisms—a global data bank of relationships between mass, elemental composition and energy content. *J Sea Res* 64: 334–340
- Brind'Amour A, Laffargue P, Morin J, Vaz S, Foveau A, Le Bris H (2014) Morphospecies and taxonomic sufficiency of benthic megafauna in scientific bottom trawl surveys. *Cont Shelf Res* 72: 1–9
- Brown CJ, Broadley A, Adame MF, Branch TA, Turschwell MP, Connolly RM (2019) The assessment of fishery status depends on fish habitats. *Fish Fish* 20: 1–14
- Brown EJ, Vasconcelos RP, Wennhage H, Bergström U and others (2018) Conflicts in the coastal zone: human impacts on commercially important fish species utilizing coastal habitat. *ICES J Mar Sci* 75: 1203–1213
- Burd BJ, Macdonald TA, van Roodselaar A (2012) Towards predicting basin-wide invertebrate organic biomass and production in marine sediments from a coastal sea. *PLOS ONE* 7: e40295
- Cebrian J (2015) Energy flows in ecosystems. *Science* 349: 1053–1054
- Christensen V, Walters CJ, Pauly D (2005) *Ecopath with Ecosim: a user's guide*. Fisheries Centre, University of British Columbia, Vancouver
- Ciotti BJ, Targett TE, Nash RDM, Geffen AJ (2014) Growth dynamics of European plaice *Pleuronectes platessa* L. in nursery areas: a review. *J Sea Res* 90: 64–82
- Collie JS (1987) Food consumption by yellowtail flounder in relation to production of its benthic prey. *Mar Ecol Prog Ser* 36: 205–213
- Craig JK, Rice JA, Crowder LB, Nadeau DA (2007) Density-dependent growth and mortality in an estuary-dependent fish: an experimental approach with juvenile spot *Leiostomus xanthurus*. *Mar Ecol Prog Ser* 343: 251–262
- de Groot AD (1969) *Methodology: foundations of inference and research in the behavioral sciences*. Mouton Publishers, Paris
- De Vlas J (1979) Annual food intake by plaice and flounder in a tidal flat area in the Dutch Wadden Sea, with special reference to consumption of regenerating parts of macrobenthic prey. *Neth J Sea Res* 13: 117–153
- Desaunay Y, Perddou JB, Beillois P (1981) Etude des nurseries de poissons du littoral de la Loire-Atlantique. *Science et Pêche* 319: 1–23
- Dorel D, Koutsikopoulos C, Desaunay Y, Marchand J (1991) Seasonal distribution of young sole (*Solea solea* (L.)) in the nursery ground of the bay of Vilaine (northern Bay of Biscay). *Neth J Sea Res* 27: 297–306
- Edwards RRC, Steele JH, Trevallion A (1970) The ecology of 0-group plaice and common dabs in Loch Ewe. III. Prey-predator experiments with plaice. *J Exp Mar Biol Ecol* 4: 156–173
- Elliott M, Hemingway K (2002) *Fishes in estuaries*. Blackwell Science, Oxford
- Elliott JM, Persson L (1978) The estimation of daily rates of food consumption for fish. *J Anim Ecol* 47: 977–991
- Elliott M, Quintino V (2007) The estuarine quality paradox, environmental homeostasis and the difficulty of detecting anthropogenic stress in naturally stressed areas. *Mar Pollut Bull* 54: 640–645
- Evans S (1983) Production, predation and food niche segregation in a marine shallow soft-bottom community. *Mar Ecol Prog Ser* 10: 147–157
- Fiksen Ø, Jørgensen C (2011) Model of optimal behaviour in fish larvae predicts that food availability determines survival, but not growth. *Mar Ecol Prog Ser* 432: 207–219
- Fonds M, Cronie R, Vethaak AD, Van Der Puyl P (1992) Metabolism, food consumption and growth of plaice (*Pleuronectes platessa*) and flounder (*Platichthys flesus*) in relation to fish size and temperature. *Neth J Sea Res* 29: 127–143
- Freitas V, Kooijman SALM, van der Veer HW (2012) Latitudinal trends in habitat quality of shallow-water flatfish nurseries. *Mar Ecol Prog Ser* 471: 203–214

- Fulford RS, Peterson MS, Grammer PO (2011) An ecological model of the habitat mosaic in estuarine nursery areas: Part I—interaction of dispersal theory and habitat variability in describing juvenile fish distributions. *Ecol Model* 222: 3203–3215
- Gee JM, Warwick RM, Davey JT, George CL (1985) Field experiments on the role of epibenthic predators in determining prey densities in an estuarine mudflat. *Estuar Coast Shelf Sci* 21: 429–448
- Gibson R (1994) Impact of habitat quality and quantity on the recruitment of juvenile flatfishes. *Neth J Sea Res* 32: 191–206
- Gibson RN, Robb L, Wennhage H, Burrows MT (2002) Ontogenetic changes in depth distribution of juvenile flatfishes in relation to predation risk and temperature on a shallow-water nursery ground. *Mar Ecol Prog Ser* 229: 233–244
- Hamerlynck O, Hostens K (1993) Growth, feeding, production, and consumption in 0-group bib (*Trisopterus luscus* L.) and whiting (*Merlangius merlangus* L.) in a shallow coastal area of the south-west Netherlands. *ICES J Mar Sci* 50: 81–91
- Hampel H, Cattrijsse A, Elliott M (2005) Feeding habits of young predatory fishes in marsh creeks situated along the salinity gradient of the Schelde estuary, Belgium and The Netherlands. *Helgol Mar Res* 59: 151–162
- Hare JA (2014) The future of fisheries oceanography lies in the pursuit of multiple hypotheses. *ICES J Mar Sci* 71: 2343–2356
- Heath MR (2005) Changes in the structure and function of the North Sea fish foodweb, 1973–2000, and the impacts of fishing and climate. *ICES J Mar Sci* 62: 847–868
- Heithaus MR, Frid A, Wirsing AJ, Worm B (2008) Predicting ecological consequences of marine top predator declines. *Trends Ecol Evol* 23: 202–210
- Hidalgo F, Alliot E, Thebault H (1987) Influence of water temperature on food intake, food efficiency and gross composition of juvenile sea bass, *Dicentrarchus labrax*. *Aquaculture* 64: 199–207
- Houde E (2008) Emerging from Hjort's shadow. *J Northwest Atl Fish Sci* 41: 53–70
- Iles TC, Beverton RJH (2000) The concentration hypothesis: the statistical evidence. *ICES J Mar Sci* 57: 216–227
- Johnson DW (2007) Habitat complexity modifies post-settlement mortality and recruitment dynamics of a marine fish. *Ecology* 88: 1716–1725
- Juanes F (2007) Role of habitat in mediating mortality during the post-settlement transition phase of temperate marine fishes. *J Fish Biol* 70: 661–677
- Jung AS, Dekker R, Germain M, Philippart CJM, Witte JIJ, van der Veer HW (2017) Long-term shifts in intertidal predator and prey communities in the Wadden Sea and consequences for food requirements and supply. *Mar Ecol Prog Ser* 579: 37–53
- Kooijman SALM (2009) Dynamic energy budget theory for metabolic organisation. Cambridge University Press, Cambridge
- Kopp D, Le Bris H, Grimaud L, Nerot C, Brind'Amour A (2013) Spatial analysis of the trophic interactions between two juvenile fish species and their preys along a coastal-estuarine gradient. *J Sea Res* 81: 40–48
- Kostecki C, Le Loc'h F, Roussel JM, Desroy N and others (2010) Dynamics of an estuarine nursery ground: the spatio-temporal relationship between the river flow and the food web of the juvenile common sole (*Solea solea*, L.) as revealed by stable isotopes analysis. *J Sea Res* 64: 54–60
- Le Bris H, Glemarec M (1995) Macrozoobenthic communities of an oxygen under-saturated coastal ecosystem: the Bay of Vilaine (southern Brittany). *Oceanol Acta* 18: 573–581
- Le Pape O, Bonhommeau S (2015) The food limitation hypothesis for juvenile marine fish. *Fish Fish* 16: 373–398
- Le Pape O, Desaunay Y, Guerault D (2003) Relationship between fluvial discharge and sole (*Solea solea*, L.) recruitment in the Bay of Biscay (France). *ICES Mar Sci Symp* 219: 241–248
- Le Pape O, Baulier L, Cloarec A, Martin J, Le Loc'h F, Désaunay Y (2007) Habitat suitability for juvenile common sole (*Solea solea*, L.) in the Bay of Biscay (France): a quantitative description using indicators based on epibenthic fauna. *J Sea Res* 57: 126–136
- Levin P, Petrik R, Malone J (1997) Interactive effects of habitat selection, food supply and predation on recruitment of an estuarine fish. *Oecologia* 112: 55–63
- Libralato S, Coll M, Tudela S, Palomera I, Pranovi F (2008) Novel index for quantification of ecosystem effects of fishing as removal of secondary production. *Mar Ecol Prog Ser* 355: 107–129
- Lindeman RL (1942) The trophic-dynamic aspect of ecology. *Ecology* 23: 399–417
- Lobry J, David V, Pasquaud S, Lepage M, Sautour B, Rochard E (2008) Diversity and stability of an estuarine trophic network. *Mar Ecol Prog Ser* 358: 13–25

- Mackinson S, Daskalov D (2007) An ecosystem model of the North Sea to support an ecosystem approach to fisheries management: description and parameterisation. *Cefas*, Lowestoft
- McLusky DS, Elliott M (2004) *The estuarine ecosystem: ecology, threats and management*. Oxford University Press, Oxford
- Metropolis N, Ulam S (1949) The Monte Carlo method. *J Am Stat Assoc* 44: 335–341
- Minto C, Myers RA, Blanchard W (2008) Survival variability and population density in fish populations. *Nature* 452: 344–347
- Monaco ME, Ulanowicz RE (1997) Comparative ecosystem trophic structure of three US mid-Atlantic estuaries. *Mar Ecol Prog Ser* 161: 239–254
- Nash RDM, Geffen AJ (2000) The influence of nursery ground processes in the determination of year-class strength in juvenile plaice *Pleuronectes platessa* L. in Port Erin Bay, Irish Sea. *J Sea Res* 44: 101–110
- Nash RDM, Geffen AJ (2012) Mortality through the early life-history of fish: What can we learn from European plaice (*Pleuronectes platessa* L.)? *J Mar Syst* 93: 58–68
- Nash RDM, Geffen AJ, Burrows MT, Gibson RN (2007) Dynamics of shallow-water juvenile flatfish nursery grounds: application of the self-thinning rule. *Mar Ecol Prog Ser* 344: 231–244
- Nicolas D, Le Loc'h F, Désaunay Y, Hamon D, Blanchet A, Le Pape O (2007) Relationships between benthic macrofauna and habitat suitability for juvenile common sole (*Solea solea*, L.) in the Vilaine estuary (Bay of Biscay, France) nursery ground. *Estuar Coast Shelf Sci* 73: 639–650
- Nicolas D, Lobry J, Le Pape O, Boët P (2010) Functional diversity in European estuaries: relating the composition of fish assemblages to the abiotic environment. *Estuar Coast Shelf Sci* 88: 329–338
- Peterson CH, Summerson HC, Thomson E, Lenihan HS and others (2000) Synthesis of linkages between benthic and fish communities as a key to protecting essential fish habitat. *Bull Mar Sci* 66: 759–774
- Pihl L (1985) Food selection and consumption of mobile epibenthic fauna in shallow marine areas. *Mar Ecol Prog Ser* 22: 169–179
- Raimonet M, Cloern JE (2017) Estuary–ocean connectivity: fast physics, slow biology. *Glob Change Biol* 23: 2345–2357
- Reiss H, Kröncke I, Ehrich S (2006) Estimating the catching efficiency of a 2-m beam trawl for sampling epifauna by removal experiments. *ICES J Mar Sci* 63: 1453–1464
- Rijnsdorp AD, Van Beek FA, Flatman S, Millner RM, Riley JD, Giret M, De Clerck R (1992) Recruitment of sole stocks, *Solea solea* (L.), in the Northeast Atlantic. *Neth J Sea Res* 29: 173–192
- Ritter C, Montagna PA, Applebaum S (2005) Short-term succession dynamics of macrobenthos in a salinity-stressed estuary. *J Exp Mar Biol Ecol* 323: 57–69
- Ross SW (2003) The relative value of different estuarine nursery areas in North Carolina for transient juvenile marine fishes. *Fish Bull* 101: 384–404
- Rybarczyk H, Elkaim B (2003) An analysis of the trophic network of a macrotidal estuary: the Seine Estuary (Eastern Channel, Normandy, France). *Estuar Coast Shelf Sci* 58: 775–791
- Rybarczyk H, Elkaim B, Ochs L, Loquet N (2003) Analysis of the trophic network of a macrotidal ecosystem: the Bay of Somme (Eastern Channel). *Estuar Coast Shelf Sci* 58: 405–421
- Ryer CH, Laurel BJ, Stoner AW (2010) Testing the shallow water refuge hypothesis in flatfish nurseries. *Mar Ecol Prog Ser* 415: 275–282
- Salen-Picard C, Darnaude AM, Arlhac D, Harmelin-Vivien ML (2002) Fluctuations of macrobenthic populations: a link between climate-driven river run-off and sole fishery yields in the Gulf of Lions. *Oecologia* 133: 380–388
- Saulnier E, Brind'Amour A, Tableau A, Rufino MM, Dauvin JC, Luczak C, Le Bris H (2019) Seasonality in coastal macrobenthic biomass and its implications for estimating secondary production using empirical models. *Limnol Oceanogr* 64: 935–949
- Schückel S, Sell AF, Kröncke I, Reiss H (2012) Diet overlap among flatfish species in the southern North Sea. *J Fish Biol* 80: 2571–2594
- Seitz RD, Wennhage H, Bergström U, Lipcius RN, Ysebaert T (2014) Ecological value of coastal habitats for commercially and ecologically important species. *ICES J Mar Sci* 71: 648–665
- Shaw M, Jenkins GP (1992) Spatial variation in feeding, prey distribution and food limitation of juvenile flounder *Rhombosolea tapirina* Günther. *J Exp Mar Biol Ecol* 165: 1–21
- Sheaves M, Baker R, Nagelkerken I, Connolly RM (2015) True value of estuarine and coastal nurseries for fish: incorporating complexity and dynamics. *Estuaries Coasts* 38: 401–414

- Sinclair ARE, Arcese P (eds) (1995) Serengeti II: dynamics, management, and conservation of an ecosystem. University of Chicago Press, Chicago, IL
- Sinclair ARE, Pech RP (1996) Density dependence, stochasticity, compensation and predator regulation. *Oikos* 75: 164–173
- Sköld M, Loo LO, Rosenberg R (1994) Production, dynamics and demography of an *Amphiura filiformis* population. *Mar Ecol Prog Ser* 103: 81–90
- Sogard SM (1997) Size-selective mortality in the juvenile stage of teleost fishes: a review. *Bull Mar Sci* 60: 1129–1157
- Tableau A, Le Bris H, Brind'Amour A (2015) Available benthic energy coefficient (ABEC): a generic tool to estimate the food profitability in coastal fish nurseries. *Mar Ecol Prog Ser* 522: 203–218
- Tableau A, Brind'Amour A, Woillez M, Le Bris H (2016) Influence of food availability on the spatial distribution of juvenile fish within soft sediment nursery habitats. *J Sea Res* 111: 76–87
- Tecchio S, Rius AT, Dauvin JC, Lobry J and others (2015) The mosaic of habitats of the Seine estuary: insights from food-web modelling and network analysis. *Ecol Model* 312: 91–101
- van der Veer HW, Witte JIJ (1993) The 'maximum growth/ optimal food condition' hypothesis: a test for 0-group plaice *Pleuronectes platessa* in the Dutch Wadden Sea. *Mar Ecol Prog Ser* 101: 81–90
- van der Veer HW, Geffen AJ, Witte JIJ (2000) Exceptionally strong year classes in plaice *Pleuronectes platessa*: Are they generated during the pelagic stage only, or also in the juvenile stage? *Mar Ecol Prog Ser* 199: 255–262
- van der Veer HW, Freitas V, Koot J, Witte JIJ, Zuur AF (2010) Food limitation in epibenthic species in temperate intertidal systems in summer: analysis of 0-group plaice *Pleuronectes platessa*. *Mar Ecol Prog Ser* 416: 215–227
- van der Veer HW, Koot J, Aarts G, Dekker R, Diderich W, Freitas V, Witte JIJ (2011) Long-term trends in juvenile flatfish indicate a dramatic reduction in nursery function of the Balgzand intertidal, Dutch Wadden Sea. *Mar Ecol Prog Ser* 434: 143–154
- van der Veer HW, Jung AS, Freitas V, Philippart CJM, Witte JIJ (2016) Possible causes for growth variability and summer growth reduction in juvenile plaice *Pleuronectes platessa* L. in the western Dutch Wadden Sea. *J Sea Res* 111: 97–106
- Vethaak AD (1992) Diseases of flounder (*Platichthys flesus* L.) in the Dutch Wadden Sea, and their relation to stress factors. *Neth J Sea Res* 29: 257–271
- Vignon M, Dierking J (2011) Prey regurgitation and stomach vacuity among groupers and snappers. *Environ Biol Fishes* 90: 361–366
- Vinagre C, Cabral HN (2008) Prey consumption by the juvenile soles, *Solea solea* and *Solea senegalensis*, in the Tagus estuary, Portugal. *Estuar Coast Shelf Sci* 78: 45–50
- Walters CJ, Juanes F (1993) Recruitment limitation as a consequence of natural selection for use of restricted feeding habitats and predation risk taking by juvenile fishes. *Can J Fish Aquat Sci* 50: 2058–2070
- Watson R, Zeller D, Pauly D (2014) Primary productivity demands of global fishing fleets. *Fish Fish* 15: 231–241
- Wetz MS, Hutchinson EA, Lunetta RS, Paerl HW, Taylor JC (2011) Severe droughts reduce estuarine primary productivity with cascading effects on higher trophic levels. *Limnol Oceanogr* 56: 627–638
- Wouters N, Cabral HN (2009) Are flatfish nursery grounds richer in benthic prey? *Estuar Coast Shelf Sci* 83: 613–620
- Zhou S, Smith AD, Knudsen EE (2015) Ending overfishing while catching more fish. *Fish Fish* 16: 716–722

Novel approach for testing the food limitation hypothesis in estuarine and coastal fish nurseries

A. Tableau*, H. Le Bris, E. Saulnier, O. Le Pape, A. Brind'Amour

Supplement S1: Statistical distribution of the uncertain parameters

The variance (V) and the mean (E) of literature data are used to define a beta distribution for ratio parameters (comprised between 0 and 1), and a gamma distribution for the strictly positive parameters. The relationships between the variance (V) and the mean (E) of data to the shape parameters of the beta distribution (α and β), and to the shape (α) and rate (β) parameters of the gamma distribution are:

beta distribution coefficients: $\alpha = \frac{E(E-E^2-V)}{V}$ (S1.1)

$$\beta = \frac{(1-E)(E-E^2-V)}{V} \quad (\text{S1.2})$$

gamma distribution coefficients: $\alpha = \frac{E^2}{V}$ (S2.1)

$$\beta = \frac{E}{V} \quad (\text{S2.2})$$

A parameter defined as a variable X whose logarithm is normally distributed (with a known mean: E and a known variance: V) follows a log-normal distribution, which is directly defined by E and V . The mean (E') of X can be expressed in function of the mean (E) and variance (V) of $\log(X)$:

mean of lognormal distribution: $E' = e^{(E+\frac{V}{2})}$ (S3)

An exception is applied to the parameter K , which is in the denominator of the consumption equation. Indeed, if K follows a beta distribution, the distribution of $1/K$ is unknown and has an unstable mean. K is thus defined with a gamma distribution. Therefore, $1/K$ is defined as an inverse gamma distribution, which is defined by the shape (α) and rate (β) parameters of the K distribution. The mean (E') of $1/K$ can be expressed in function of the mean (E) and variance (V) of K :

mean of inverse gamma distribution: $E' = \frac{E}{(E^2-V)}$ with the constraint: $V < E^2$ (S4)

Supplement S2: data collection for the Bay of Vilaine

Survey data

Macrobenthic invertebrate biomass and fish abundance

Given the mismatch in the locations of the sampling stations between the two surveys (fish and invertebrates) and the uneven spatial distribution of the sampling effort, we stratified the bay of Vilaine in three zones (Figure 1, which is a simplification of the 5 habitats defined by Kopp et al. 2013). Splitting is necessary because the sampling efforts of fish and benthic invertebrates decrease from zone 1 to zone 2, and zone 3 (respectively 3.4 stations.km⁻², 2.0 stations.km⁻², and 1.5 stations.km⁻² for fish sampling; and respectively 0.42 stations.km⁻², 0.17 stations.km⁻², and 0.17 stations.km⁻² for benthic invertebrate sampling). Therefore, for each cohort/species, the average abundance/biomass is computed in each zone and raised to the surface area to provide a total abundance/biomass. Bootstrap method is used to estimate the uncertainty on total abundance/biomass of each cohort/species.

Fish weight-length relationships

For each haul of the survey, fish are weighed by species all together, and they are all individually measured (a subsample is measured if the number exceeds 30 individuals). An observation is described as follows:

$$W_h = \sum_{n \in 1:N} a \cdot L_n^b \quad (\text{S5})$$

W_h is the weight in g. of all individuals of a single species in the haul h ; L_n in mm is the length of the n^{th} fish individual; a and b , the parameters of the relationship. These parameters are then estimated using all observations with an optimization algorithm in the software R (R Core Team 2012).

Table S1: weight-length parameters (weight in g and length in mm)

Species	a	b
<i>Dicologlossa cuneata</i>	3.028e-5	2.723
<i>Pleuronectes platessa</i>	1.369e-5	2.953
<i>Solea solea</i>	9.237e-6	2.994
<i>Mullus surmuletus</i>	7.636e-6	3.121
<i>Merlangius merlangus</i>	9.749e-6	2.972
<i>Merluccius merluccius</i>	5.454e-6	3.055
<i>Trisopterus luscus</i>	9.705e-6	3.054

Literature data

Macrobenthic invertebrates

Table S2: Summary of macrobenthic invertebrate parameters.

Parameter	Value or range of values	Uncertainty details	Source
Conversion ratio (CR)	0.672	sd=0.0111	survey
Production-to-biomass ratio (P:B)	0.222-9.723 year ⁻¹	sd range: 0.0356-1.988	(Brey 2012)
Regeneration (R)	0-0.5	Conservative approach	various sources
Energy conversion (E)	0.851-6.694 kJ.g ⁻¹	Negligible uncertainty	(Brey et al. 2010)
Accessibility (A)	Easily accessible : 1 Hardly accessible : 0.117	sd=0.0485	survey

Additional details on these parameters are provided in Tableau et al 2015 for the 94 taxa.

Fish juveniles

Table S3: Fish data compiled from the literature. * As *Merluccius merluccius* shifts to an exclusive fish diet after 150 mm, this proportion becomes 0 for larger fish.

fish	<i>Dicologlossa cuneata</i>	<i>Pleuronectes platessa</i>	<i>Solea solea</i>	<i>Solea solea</i>	<i>Mullus surmuletus</i>	<i>Merlangius merlangus</i>	<i>Merluccius merluccius</i>	<i>Trisopterus luscus</i>
age group	G1	G0	G0	G1	G0	G0	G0	G0
shape	flatfish	flatfish	flatfish	flatfish	roundfish	roundfish	roundfish	roundfish
diet	exclusive benthivore	exclusive benthivore	exclusive benthivore	exclusive benthivore	exclusive benthivore	partial benthivore	partial benthivore	partial benthivore
Ld₀ (mm)	110	70	70	110	70	70	70	70
d0	April 11	August 7	July 31	May 21	August 11	July 12	April 9	July 9
G (mm.day⁻¹)	0.38	0.51	0.60	0.60	0.77	0.79	0.67	0.83
Z (day⁻¹)	0.0179	0.0171	0.0179	0.0179	0.0103	0.00760	0.00765	0.00770
q	0.257	0.380	0.257	0.257	0.500	0.500	0.500	0.500
DC	1	1	1	1	1	0.5	0.27*	0.75
E (kJ.g⁻¹)	5.78	5.66	5.78	5.78	6.88	6.24	6.24	6.24
E sd	0.618	0.802	0.618	0.618	1.69	0.805	0.805	0.805
K	0.197	0.316	0.197	0.197	0.312	0.385	0.385	0.385
K sd	0.0675	0.0792	0.0675	0.0675	0.0969	0.0720	0.0720	0.0720

Growth period and growth rate

The main growth period of fish juveniles in temperate ecosystems is ranging from spring to fall and is very consistent over many fish species (Forest 1975, Gordon 1977, Desaunay et al. 1981, Van Der Veer et al. 1990, Hamerlynck & Hostens 1993, Shi et al. 1997, Amara 2004, Otxotorena et al. 2010). This is explained by lower food intake during winter (Zanuy & Carrillo 1985). By conservative approach, we focus only on the food intake during the main growth period in summer. As a growth decrease is generally observed during fall, we set the end of the growth period in our model on October 1 (early fall). For G0 fish, the beginning of the growth period is deduced from the growth rate and the fish size from which they eat on macrobenthic invertebrates (set at 70 mm, see Tableau et al. 2016). For G1 *Solea solea*, the length at the beginning of the growth period is associated to the length of G0 at the end of the growth period (11 cm), this is supported by the observations in Desaunay et al. (1981). As no G0 *Dicologlossa cuneata* are observed, we use the same length for G1 *Dicologlossa cuneata*. This assumption is supported by the observations in Forest (Forest 1975).

The growth rates can vary over time and among the nursery habitats (Ciotti et al. 2014). The general strategy was to use data corresponding to the closest habitats of the bay of Vilaine when no local data were available. When length time series were available, the match with the fish length observed in the survey was checked (Forest 1975, Gordon 1977, Desaunay et al. 1981, Van Der Veer et al. 1990, Hamerlynck & Hostens 1993, Shi et al. 1997, Amara 2004, Otxotorena et al. 2010). The conservative approach for the growth period limits the potential overestimation of the exploitation efficiency.

Table S4: growth data collection (G0 for young of the year and G1 for one year old)

Species	Age group	Growth rate (mm.day ⁻¹)	Source	Comment
<i>Dicologlossa cuneata</i>	G1	0.38	Forest 1975	graphic analysis on local data
<i>Pleuronectes platessa</i>	G0	0.51	Amara 2004	local data
<i>Solea solea</i>	G0	0.60	Marchand 1991	local data
<i>Solea solea</i>	G1	0.60	Desaunay et al. 1981, Marchand 1991	local data & graphic analysis on local data
<i>Mullus surmuletus</i>	G0	0.77	N'Da and Deniel 2005	data analysis on local data
<i>Merlangius merlangus</i>	G0	0.79	Hamerlynck and Hostens 1993	graphic analysis on external data - good match between lengths at the same period
<i>Merluccius merluccius</i>	G0	0.67	Kacher and Amara 2005, Otxotorena et al. 2010	average on two local data
<i>Trisopterus luscus</i>	G0	0.83	Hamerlynck and Hostens 1993	graphic analysis on external data - good match between lengths at the same period

Mortality rate

The mortality rates vary among nursery habitats and over time (Iles & Beverton 1991). In the exploitation efficiency analysis, emigration and natural mortality can be confounded as it results in a decrease of food consumption anyway. Mortality rates are available only for two flatfish species and are used for the others. A single data is available for three roundfish species, the average of the two Gadiformes species (*Merlangius merlangus* and *Trisopterus luscus*) is used for *Merluccius merluccius*:

Pleuronectes platessa (51 data): mean = 0.0171 nb.day⁻¹ & sd = 0.0088 (Iles & Beverton 1991, Modin & Pihl 1994, Nash & Geffen 2012)

Limanda limanda (9 data): mean = 0.0187 nb.day⁻¹ & sd = 0.0103 (Iles & Beverton 1991)

Mullus surmuletus (1 data): mean = 0.0103 (N'Da & Deniel 2005)

Merlangius merlangus (1 data): mean = 0.0076 (Hamerlynck & Hostens 1993)

Trisopterus luscus (1 data): mean = 0.0077 (Hamerlynck & Hostens 1993)

As the survey is done around the middle of the growth period (August 22), errors in mortality rates have small consequences: an overestimated mortality would result in a higher consumption estimate on the time period before the survey, but a lower one on the period after the survey. As we want to estimate the exploitation on the full time period, the lack of accuracy of this parameter is acceptable.

Catch efficiency

Catch efficiency data have been estimated in several studies on various flatfish species including *Pleuronectes platessa* and *Solea solea*. The catch efficiencies are most likely overestimated in these studies given the experimental protocols (e.g Kuipers 1975). Consequently, fish abundances are probably underestimated, which fits with the conservative approach. First, available data have been compiled per study and then averaged over the different studies. *Solea solea* value has been attributed to *Dicologlossa cuneata* given their close phylogeny (*Soleidae* family). Data on roundfish species are scarcer but conservative data are suggested for *Merlangius merlangius* and *Trisopterus luscus*. This data has been associated to the roundfish *Mullus surmulutus* and *Merluccius merluccius*.

Table S5: catch efficiency data

Species	q	Trawl	Source
<i>Pleuronectes platessa</i>	50.6 %	2 m beam trawl	(Rogers & Lockwood 1989)
<i>Pleuronectes platessa</i>	28.4 %	2 m beam trawl	(Kuipers 1975)
<i>Pleuronectes platessa</i>	39.5 %	2 m beam trawl	(Kuipers 1975)
<i>Pleuronectes platessa</i>	33.5 %	2 m beam trawl	(Edwards & Steele 1968)
<i>Solea solea</i>	25.7 %	2 m beam trawl	(Rogers & Lockwood 1989)
<i>Merlangius merlangus</i>	50.0 %	3 m beam trawl	(Hamerlynck & Hostens 1993)
<i>Trisopterus luscus</i>	50.0 %	3 m beam trawl	(Hamerlynck & Hostens 1993)

Diet

The first age groups of the three flatfish species are exclusive benthivorous species (Amara et al. 2001, Le Loc'h 2004). *Mullus surmulletus* is also an exclusive benthivore (N'da 1992). Based on the gravimetric proportion of benthic invertebrate prey observed in gut contents of G0 *Merlangius merlangus* and *Trisopterus luscus* (Hamerlynck & Hostens 1993), the estimated proportions are 50 % and 75 % respectively. Based on Mahe et al. (2007), the proportions in the *Merluccius merluccius* diet of benthic invertebrates are averaged over three regions (27 %). As G0 *Merluccius merluccius* diet shifts from a mix of fish and benthic invertebrates to only fish around 15mm, the proportion is set to 0 after 150 mm (Le Loc'h 2004, Mahe et al. 2007).

Energy density

An investigation has been done on all articles of the journals “Aquaculture”, “Aquaculture nutrition” and “Aquaculture research”. A complementary research in all journals has been done with the keywords “Body composition” “energy density”. The body composition is often described as a proportion of proteins, lipids, glycogen, and water. Energy conversion coefficients have been compiled from different studies (Brett & Groves 1979, Brafield 1985, Cho & Kaushik 1990, Jobling 1993).

Proteins: 23.7kJ.g⁻¹

Lipids: 36.3kJ.g⁻¹

Glycogen: 17.2kJ.g⁻¹

Water: 0kJ.g⁻¹

As adult body composition varies with the reproduction season, only juvenile data are used. Pelagic fish data are rejected as this kind of fish has a higher fat concentration. Data corresponding to fish fed with high fat food are rejected. All data referring only to the flesh or the liver were also rejected. Among the 186 data collected, 62 referring to the fish taxa *Gadidae*, *Perciformes*, *Pleuronectidae*, and *Soleidae* were used. The average and the standard deviation of the corresponding taxa are attributed to each species. 25 references are used (Pandian 1970, Holdway & Beamish 1984, Smith et al. 1986, Hidalgo et al. 1987, Costopoulos & Fonds 1989, Marais 1990, Imsland et al. 2000, Lee et al. 2003, Person-Le Ruyet et al. 2004, Dias et al. 2004, Rosenlund et al. 2004, Tibbetts et al. 2005, Rema et al. 2008, Borges et al. 2009, Silva et al. 2010, Ding et al. 2010, Gatta et al. 2011, Li et al. 2012, Marinho et al. 2014, Guerreiro et al. 2015, Bonvini et al. 2015, López et al. 2016, Zhang et al. 2016, Li et al. 2017, Salas-Leiton et al. 2017).

Gross conversion efficiency (growth-to-food intake ratio)

An investigation has been done on all articles of the journals “Aquaculture”, “Aquaculture nutrition” and “Aquaculture research”. A complementary research has been done in all journals with the keywords “growth efficiency energy”, “gross growth efficiency energy”, “gross conversion efficiency energy”, “food conversion efficiency energy”, “feed conversion efficiency energy”. Data collected in winter are rejected as lower temperature gives lower conversion efficiency (e.g. Zanuy & Carrillo 1985). When conversion efficiency data are estimated on a temperature gradient, we take only values around the optimal temperature. When data refer to weight conversion efficiency, we convert the production and the consumption in energy when possible. Gross conversion efficiencies are higher for young fish as no energy is allocated to reproduction. Thus, we use only gross conversion efficiencies referring to fish juveniles (Mateo 2007). We used 189 data referring to the fish taxa *Gadidae*, *Perciformes*, *Pleuronectidae*, and *Soleidae*. As these data mostly come from aquaculture studies, they are expected to be a bit higher than the gross conversion efficiency occurring in natural environment. The exploitation efficiencies are consequently likely underestimated. The average and the standard deviation of the corresponding taxa are attributed to each species. 31 references are used (Pandian 1970, Edwards et al. 1972, Marais & Kissil 1979, Hidalgo et al. 1987, Fonds et al. 1992, Hamerlynck & Hostens 1993, Lankford & Targett 1994, Buckel et al. 1995, Björnsson & Tryggvadóttir 1996, Deacon & Hecht 1999, Imsland et al. 2000, Kim & Lall 2001, Kim et al. 2001, Peck et al. 2003, Person-Le Ruyet et al. 2004, Dias et al. 2004, Rosenlund et al. 2004, Tibbetts et al. 2005, Mateo 2007, Rema et al. 2008, Borges et al. 2009, Silva et al. 2010, Ding et al. 2010, Gatta et al. 2011, Guerreiro et al. 2012, Li et al. 2012, Bonvini et al. 2015, López et al. 2016, Zhang et al. 2016, Li et al. 2017, Salas-Leiton et al. 2017).

References

- Amara R (2004) 0-group flatfish growth conditions on a nursery ground (Bay of Canche, Eastern English Channel). *Hydrobiologia* 518:23–32
- Amara R, Laffargue P, Dewarumez JM, Maryniak C, Lagardère F, Luzac C (2001) Feeding ecology and growth of 0-group flatfish (sole, dab and plaice) on a nursery ground (Southern Bight of the North Sea). *J Fish Biol* 58:788–803
- Björnsson B, Tryggvadóttir SV (1996) Effects of size on optimal temperature for growth and growth efficiency of immature Atlantic halibut (*Hippoglossus hippoglossus* L.). *Aquaculture* 142:33–42
- Bonvini E, Parma L, Mandrioli L, Sirri R, Brachelente C, Mongile F, Gatta PP, Bonaldo A (2015) Feeding common sole (*Solea solea*) juveniles with increasing dietary lipid levels affects growth, feed utilization and gut health. *Aquaculture* 449:87–93
- Borges P, Oliveira B, Casal S, Dias J, Conceição L, Valente LMP (2009) Dietary lipid level affects growth performance and nutrient utilisation of Senegalese sole (*Solea senegalensis*) juveniles. *Br J Nutr* 102:1007–1014
- Brafield AE (1985) Laboratory Studies of Energy Budgets. In: Tytler P, Calow P (eds) *Fish Energetics*. Springer Netherlands, p 257–281
- Brett JR, Groves TDD (1979) Physiological energetics. In: *Fish Physiology*. Academic Press, New York, p 280–352
- Brey T (2012) A multi-parameter artificial neural network model to estimate macrobenthic invertebrate productivity and production. *Limnol Oceanogr-Methods* 10:581–589
- Brey T, Mueller-Wiegmann C, Zittier ZMC, Hagen W (2010) Body composition in aquatic organisms - A global data bank of relationships between mass, elemental composition and energy content. *J Sea Res* 64:334–340
- Buckel JA, Steinberg ND, Conover DO (1995) Effects of temperature, salinity, and fish size on growth and consumption of juvenile bluefish. *J Fish Biol* 47:696–706
- Cho CY, Kaushik SJ (1990) Nutritional energetics in fish: energy and protein utilization in rainbow trout (*Salmo gairdneri*). In: *Aspects of food production, consumption and energy values*. Karger Publishers, p 132–172
- Ciotti BJ, Targett TE, Nash RDM, Geffen AJ (2014) Growth dynamics of European plaice *Pleuronectes platessa* L. in nursery areas: A review. *J Sea Res* 90:64–82
- Costopoulos CG, Fonds M (1989) Proximate body composition and energy content of plaice (*Pleuronectes platessa*) in relation to the condition factor. *Neth J Sea Res* 24:45–55
- Deacon N, Hecht T (1999) The effect of reduced salinity on growth, food conversion and protein efficiency ratio in juvenile spotted grunter, *Pomadourys commersonnii* (Lacépède) (Teleostei: Haemulidae). *Aquac Res* 30:13–20

- Desaunay Y, Perddou JB, Beillois P (1981) Etude des nurseries de poissons du littoral de la Loire-Atlantique. *Sci Pêche* 319:1–23
- Dias J, Rueda-Jasso R, Panserat S, Conceição LEC da, Gomes EF, Dinis MT (2004) Effect of dietary carbohydrate-to-lipid ratios on growth, lipid deposition and metabolic hepatic enzymes in juvenile Senegalese sole (*Solea senegalensis*, Kaup). *Aquac Res* 35:1122–1130
- Ding L, Zhang L, Wang J, Ma J, Meng X, Duan P, Sun L, Sun Y (2010) Effect of dietary lipid level on the growth performance, feed utilization, body composition and blood chemistry of juvenile starry flounder (*Platichthys stellatus*). *Aquac Res* 41:1470–1478
- Edwards RRC, Finlayson DM, Steele JH (1972) An experimental study of the oxygen consumption, growth, and metabolism of the cod (*Gadus Morhua* L.). *J Exp Mar Biol Ecol* 8:299–309
- Edwards R, Steele JH (1968) The ecology of 0-group plaice and common dabs at Loch Ewe I. Population and food. *J Exp Mar Biol Ecol* 2:215–238
- Fonds M, Cronie R, Vethaak AD, Van Der Puyl P (1992) Metabolism, food consumption and growth of plaice (*Pleuronectes platessa*) and flounder (*Platichthys flesus*) in relation to fish size and temperature. *Neth J Sea Res* 29:127–143
- Forest A (1975) Le ceteau *Dicologlossa cunetata* (Moreau), sa biologie et sa peche dans le sud du golfe de Gascogne. *Rev Trav Inst Pêch Marit* 39:5–62
- Gatta PP, Parma L, Guarniero I, Mandrioli L, Sirri R, Fontanillas R, Bonaldo A (2011) Growth, feed utilization and liver histology of juvenile common sole (*Solea solea* L.) fed isoenergetic diets with increasing protein levels. *Aquac Res* 42:313–321
- Gordon JDM (1977) The fish populations in inshore waters of the West Coast of Scotland. The distribution, abundance and growth of the whiting (*Merlangius merlangus* L.). *J Fish Biol* 10:587–596
- Guerreiro I, Enes P, Merrifield D, Davies S, Oliva-Teles A (2015) Effects of short-chain fructooligosaccharides on growth performance and hepatic intermediary metabolism in turbot (*Scophthalmus maximus*) reared at winter and summer temperatures. *Aquac Nutr* 21:433–443
- Guerreiro I, Peres H, Castro-Cunha M, Oliva-Teles A (2012) Effect of temperature and dietary protein/lipid ratio on growth performance and nutrient utilization of juvenile Senegalese sole (*Solea senegalensis*). *Aquac Nutr* 18:98–106
- Hamerlynck O, Hostens K (1993) Growth, feeding, production, and consumption in 0-group bib (*Trisopterus luscus* L.) and whiting (*Merlangius merlangus* L.) in a shallow coastal area of the south-west Netherlands. *ICES J Mar Sci J Cons* 50:81–91
- Hidalgo F, Alliot E, Thebault H (1987) Influence of water temperature on food intake, food efficiency and gross composition of juvenile sea bass, *Dicentrarchus labrax*. *Aquaculture* 64:199–207
- Holdway DA, Beamish FWH (1984) Specific growth rate and proximate body composition of Atlantic cod (*Gadus morhua* L.). *J Exp Mar Biol Ecol* 81:147–170

- Iles TC, Beverton RJH (1991) Mortality rates of 0-group plaice (*Pleuronectes platessa* L.), dab (*Limanda limanda* L.) and turbot (*Scophthalmus maximus* L.) in European waters: I. Statistical analysis of the data and estimation of parameters. *Neth J Sea Res* 27:217–235
- Imsland AK, Jonassen TM, Stefansson SO, Kadowaki S, Berntssen MHG (2000) Intraspecific Differences in Physiological Efficiency of Juvenile Atlantic Halibut *Hippoglossus hippoglossus* L. *J World Aquac Soc* 31:285–296
- Jobling M (1993) Bioenergetics: feed intake and energy partitioning. In: Rankin JC, Jensen FB (eds) *Fish Ecophysiology*. Springer Netherlands, p 1–44
- Kacher M, Amara R (2005) Distribution and growth of 0-group European hake in the Bay of Biscay and Celtic Sea: a spatial and inter-annual analyses. *Fish Res* 71:373–378
- Kim J-D, Lall SP (2001) Effects of dietary protein level on growth and utilization of protein and energy by juvenile haddock (*Melanogrammus aeglefinus*). *Aquaculture* 195:311–319
- Kim JD, Lall SP, Milley JE (2001) Dietary protein requirements of juvenile haddock (*Melanogrammus aeglefinus* L.). *Aquac Res* 32:1–7
- Kopp D, Le Bris H, Grimaud L, Nerot C, Brind'Amour A (2013) Spatial analysis of the trophic interactions between two juvenile fish species and their preys along a coastal-estuarine gradient. *J Sea Res* 81:40–48
- Kuipers B (1975) On the efficiency of a two-metre beam trawl for juvenile plaice (*Pleuronectes Platessa*). *Neth J Sea Res* 9:69–85
- Lankford TE, Targett TE (1994) Suitability of estuarine nursery zones for juvenile weakfish (*Cynoscion regalis*): effects of temperature and salinity on feeding, growth and survival. *Mar Biol* 119:611–620
- Le Loc'h F (2004) Structure, functioning and evolution of exploited soft sediment benthic communities of the North Bay of Biscay continental shelf. Université de Bretagne occidentale - Brest
- Lee S-M, Lee JH, Kim K-D (2003) Effect of dietary essential fatty acids on growth, body composition and blood chemistry of juvenile starry flounder (*Platichthys stellatus*). *Aquaculture* 225:269–281
- Li Y, Ai Q, Mai K, Xu W, Cheng Z (2012) Effects of the partial substitution of dietary fish meal by two types of soybean meals on the growth performance of juvenile Japanese seabass, *Lateolabrax japonicus* (Cuvier 1828). *Aquac Res* 43:458–466
- Li W, Wen X, Huang Y, Zhao J, Li S, Zhu D (2017) Effects of varying protein and lipid levels and protein-to-energy ratios on growth, feed utilization and body composition in juvenile *Nibea diacanthus*. *Aquac Nutr*
- López LM, Olmos Soto J, Trejo Escamilla I, Flores Ibarra M, Ochoa L, Drawbridge M, Peres H (2016) Evaluation of carbohydrate-to-lipid ratio in diets supplemented with *Bacillus subtilis* probiotic

- strain on growth performance, body composition and digestibility in juvenile white seabass (*Atractoscion nobilis*, Ayres 1860). *Aquac Res* 47:1864–1873
- Mahe K, Amara R, Bryckaert T, Kacher M, Brylinski JM (2007) Ontogenetic and spatial variation in the diet of hake (*Merluccius merluccius*) in the Bay of Biscay and the Celtic Sea. *ICES J Mar Sci J Cons* 64:1210–1219
- Marais JFK (1990) Body composition of ten marine migrants and one freshwater fish species caught in estuaries of the Eastern Cape, South Africa. *South Afr J Mar Sci* 9:135–140
- Marais JFK, Kissil GW (1979) The influence of energy level on the feed intake, growth, food conversion and body composition of *Sparus aurata*. *Aquaculture* 17:203–219
- Marchand J (1991) The influence of environmental conditions on settlement, distribution and growth of 0-group sole (*Solea solea* (L.)) in a macrotidal estuary (Vilaine, France). *Neth J Sea Res* 27:307–316
- Marinho G, Peres H, Carvalho AP (2014) Effect of feeding time on dietary protein utilization and growth of juvenile Senegalese sole (*Solea senegalensis*). *Aquac Res* 45:828–833
- Mateo I (2007) A bioenergetics based comparison of growth conversion efficiency of Atlantic cod on Georges Bank and in the Gulf of Maine. *J Northwest Atl Fish Sci*
- Modin J, Pihl L (1994) Differences in growth and mortality of juvenile plaice, *Pleuronectes platessa* L., following normal and extremely high settlement. *Neth J Sea Res* 32:331–341
- Nash RDM, Geffen AJ (2012) Mortality through the early life-history of fish: What can we learn from European plaice (*Pleuronectes platessa* L.)? *J Mar Syst* 93:58–68
- N'da K (1992) Régime alimentaire du rouget de roche *Mullus surmuletus* (Mullidae) dans le nord du Golfe de Gascogne. *Cybium* 16:159–167
- N'Da K, Deniel C (2005) Croissance des juvéniles du rouget de roche (*Mullus surmuletus*) dans le nord du golfe de Gascogne. *Cybium* 29:175–178
- Otxotorena U, Díez G, Abechuco EL de, Santurtún M, Lucio P (2010) Estimation of age and growth of juvenile hakes (*Merluccius merluccius* Linnaeus, 1758) of the Bay of Biscay and Great Sole by means of the analysis of macro and microstructure of the otoliths. *Fish Res* 106:337–343
- Pandian TJ (1970) Intake and conversion of food in the fish *Limanda limanda* exposed to different temperatures. *Mar Biol* 5:1–17
- Peck MA, Buckley L, Caldarone E, Bengtson D (2003) Effects of Food Consumption and Temperature on Growth Rate and Biochemical-Based Indicators of Growth in Early Juvenile Atlantic Cod *Gadus morhua* and Haddock *Melanogrammus aeglefinus*. *Fish Anim Vet Sci Fac Publ*
- Person-Le Ruyet J, Mahé K, Le Bayon N, Le Delliou H (2004) Effects of temperature on growth and metabolism in a Mediterranean population of European sea bass, *Dicentrarchus labrax*. *Aquaculture* 237:269–280

- R Core Team (2012) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria
- Rema P, Conceição L e. c., Evers F, Castro-Cunha M, Dinis M t., Dias J (2008) Optimal dietary protein levels in juvenile Senegalese sole (*Solea senegalensis*). *Aquac Nutr* 14:263–269
- Rogers SI, Lockwood SJ (1989) Observations on the capture efficiency of a two-metre beam trawl for juvenile flatfish. *Neth J Sea Res* 23:347–352
- Rosenlund G, Karlsen Ø, Tveit K, Mangor-Jensen A, Hemre G-I (2004) Effect of feed composition and feeding frequency on growth, feed utilization and nutrient retention in juvenile Atlantic cod, *Gadus morhua* L. *Aquac Nutr* 10:371–378
- Salas-Leiton E, Conde-Sieira M, Pelusio N, Marques A, Maia M r. g., Soengas J l., Valente L m. p. (2017) Dietary protein/carbohydrate ratio in low-lipid diets for Senegalese sole (*Solea senegalensis*, Kaup 1858) juveniles. Influence on growth performance, nutrient utilization and flesh quality. *Aquac Nutr*:n/a-n/a
- Shi YB, Gunderson DR, Sullivan PJ (1997) Growth and survival of 0(+) English sole, *Pleuronectes vetulus*, in estuaries and adjacent nearshore waters off Washington. *Fish Bull* 95:161–173
- Silva JMG, Espe M, Conceição LEC, Dias J, Costas B, Valente LMP (2010) Feed intake and growth performance of Senegalese sole (*Solea senegalensis* Kaup, 1858) fed diets with partial replacement of fish meal with plant proteins. *Aquac Res* 41:e20–e30
- Smith RL, Paul AJ, Paul JM (1986) Effect of food intake and temperature on growth and conversion efficiency of juvenile walleye pollock (*Theragra chalcogramma* (Pallas)): a laboratory study. *ICES J Mar Sci* 42:241–253
- Tableau A, Brind'Amour A, Woillez M, Le Bris H (2016) Influence of food availability on the spatial distribution of juvenile fish within soft sediment nursery habitats. *J Sea Res* 111:76–87
- Tibbetts S m., Lall S p., Milley J e. (2005) Effects of dietary protein and lipid levels and DP DE–1 ratio on growth, feed utilization and hepatosomatic index of juvenile haddock, *Melanogrammus aeglefinus* L. *Aquac Nutr* 11:67–75
- Van Der Veer HW, Creutzberg F, Dapper R, Duineveld GCA, Fonds M, Kuipers BR, Van Noort GJ, Witte JIJ (1990) On the ecology of the dragonet *Callionymus lyra* L. in the southern North Sea. *Neth J Sea Res* 26:139–150
- Zanuy S, Carrillo M (1985) Annual cycles of growth, feeding rate, gross conversion efficiency and hematocrit levels of sea bass (*Dicentrarchus labrax* L.) adapted to two different osmotic media. *Aquaculture* 44:11–25
- Zhang C -x., Huang F, Li J, Wang L, Song K, Mai K -s. (2016) Interactive effects of dietary magnesium and vitamin E on growth performance, body composition, blood parameters and antioxidant status in Japanese seabass (*Lateolabrax japonicus*) fed oxidized oil. *Aquac Nutr* 22:708–722