



The use and performance of survey-based pre-recruit abundance indices for possible inclusion in stock assessments of coastal-dependent species

Olivier Le Pape, Youen Vermard, Jérôme Guitton, Elliot Brown, Karen van de Wolfshaar, Romuald Lipcius, Josianne Støttrup, Kenneth Rose

► To cite this version:

Olivier Le Pape, Youen Vermard, Jérôme Guitton, Elliot Brown, Karen van de Wolfshaar, et al.. The use and performance of survey-based pre-recruit abundance indices for possible inclusion in stock assessments of coastal-dependent species. ICES Journal of Marine Science, 2020, 77 (5), pp.1953-1965. 10.1093/icesjms/fsaa051 . hal-03135804

HAL Id: hal-03135804

<https://institut-agro-rennes-angers.hal.science/hal-03135804>

Submitted on 14 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The use and performance of survey-based pre-recruit abundance indices for possible inclusion in stock assessments of coastal-dependent species

Le Pape Olivier ^{1,*}, Vermard Youen ², Guitton Jerome ¹, Brown Elliot J. ³, Van De Wolfshaar Karen E. ⁴, Lipcius Romuald N. ⁵, Stottrup Josianne G. ³, Rose Kenneth A. ⁶

¹ Agrocampus Ouest, ESE, INRAE, Ecol & Ecosyst Hlth, F-35042 Rennes, France.

² IFREMER, Ecol & Models Fisheries Sci, EMH, Plouzane, France.

³ Tech Univ Denmark, Natl Inst Aquat Resources DTU Aqua, Kemitorvet, Bldg 202, DK-2800 Lyngby, Denmark.

⁴ Wageningen Marine Res Ecol Dynam Grp, NL-1970 AB IJmuiden, Netherlands.

⁵ Virginia Inst Marine Sci, William & Mary, POB 1346, Gloucester Point, VA 23062 USA.

⁶ Univ Maryland, Ctr Environm Sci, Horn Point Lab, POB 775, Cambridge, MD 21613 USA.

* Corresponding author : Olivier Le Pape, email address : olivier.le.pape@agrocampus-ouest.fr

Abstract :

We reviewed the use of survey-based pre-recruit abundance indices in short-term recruitment forecasts for fish species relying on coastal habitats at the juvenile stage and that are assessed by ICES. We collated information from stock assessment reports and from a questionnaire filled out by the stock assessors. Among the 78 stocks with juvenile coastal dependence, 49 use short-term forecasts in stock assessment. Survey-based pre-recruit abundance indices were available for 35 of these stocks, but only 14 were used to forecast recruitment. The questionnaire indicated that the limited use of survey-based pre-recruit abundance indices was primarily due to sampling inefficiency, which may preclude reliable recruitment estimates. The sampling is inefficient because the juvenile coastal distribution is outside the geographical area covered by large-scale surveys or targeted coastal surveys are conducted on limited spatial and temporal scales. However, our analysis of the relationship between survey-based pre-recruit indices and assessment-generated recruitment indices revealed that survey-based pre-recruit abundance indices were sufficiently accurate to provide useful information for predicting future recruitment. We recommend expansion of the use of survey-based indices of pre-recruit abundance in stock assessment and recruitment forecasting, and consideration of how to include juveniles in ongoing and future surveys.

Keywords : coastal nursery, forecast, juvenile habitat, recruitment, stock assessment, survey

41 **Introduction**

42 Recruitment variability of many marine and coastal fish species is the main driver of
43 fluctuations in population abundance and critically depends on the highly variable mortality
44 rates of early life stages (Levin and Stunz, 2005; Juanes, 2007; Archambault et al., 2014).
45 Forecasting future recruitment has long been a focus of fisheries management (Hilborn and
46 Walters, 1992; Needle, 2001) and continues to be an essential part of evaluating fishery
47 management strategies (Kimoto et al., 2007; Stige et al., 2013; Punt, 2019). Stochastic
48 processes that occur at the egg and larval stages generate high mortality rates (typically 99.9%
49 for eggs and larvae; Le Pape and Bonhommeau, 2015), which can also be density-dependent
50 and can vary greatly from year-to-year, thereby generating large fluctuations in recruitment

(Houde, 2008; Cury et al., 2014; Szuwalski et al., 2015). Accordingly, egg and larval abundances estimated from ichthyoplankton surveys are often poorly correlated to future recruitment success. In contrast, after a “critical” stage or size (Cowan et al., 2000; Dingsor et al., 2007; Houde, 2008), juvenile fish experience considerably lower and more consistent mortality rates than eggs and larvae. Abundance, whether absolute or relative (index), can be estimated during the juvenile stage for many species (Le Pape and Bonhommeau, 2015), without major discrepancies arising from the highly variable mortality rates typical of earlier life stages. In stock assessment, pre-recruitment is considered the life stage after the transition from the highly variable early stages (eggs, larvae, and often early juveniles) to when natural mortality is largely stable (Lorenzen and Camp, 2019) but before individuals fully join the adult stock. Survey-based pre-recruit abundance indices could therefore provide reliable information on recruitment and future year-class strength (Helle et al., 2000; Zhang et al., 2010; Stige et al., 2013).

Indices estimating pre-recruit abundance can provide projections of recruitment and can inform fisheries management, especially for stocks whose exploitation is highly dependent on the juvenile stage. Such stocks depend on recruitment for determining harvest, due either to their biology (short-lived species, like small pelagics) or because high exploitation rates reduce the age of the fish harvested. For example, high exploitation rates of Atlantic cod *Gadus morhua* in the North Sea during the last five years (2012 to 2016) of the assessment resulted in immature fish constituting an average of 71% of the international landings in number (ICES, 2017c).

Coastal zones are biologically productive areas that serve as juvenile habitat for numerous marine species (Beck et al., 2001). For example, considering the species for which ICES provides advice, one-third are dependent on coastal habitats during their juvenile stage (Seitz et al., 2014), and these species account for 66% of the total landings of ICES-evaluated stocks (Brown et al., 2018a). Scientific surveys at the population scale are usually designed to estimate

density and age-structure of post-recruited fish. Many surveys focus on post-recruitment fish for specific management purposes and therefore are not designed nor appropriate for estimating pre-recruit abundance. Additionally, such post-recruitment surveys most often do not provide adequate coverage of coastal habitat on which juveniles rely (Ralph and Lipcius, 2014). When juveniles aggregate in coastal areas, survey designs that cover suitable shallow coastal habitats are required to produce reliable estimates of pre-recruit density. The timing within the year of the surveys is also important to give sufficient time for the recruits to settle in the juvenile habitats and to pass the early juveniles stages that incur highly variable survival (van der Veer, 1986; Wennhage, 2002; Nash et al., 2007). Surveys designed for other purposes may not cover the time period that is optimal for estimating recruitment from pre-recruits. Even when the surveys focus on juveniles before recruitment, they tend to be spatially localized, thereby creating challenges to extrapolate the results to the broader spatial domain of the managed stock. A valid reason for why surveys are not used to generate pre-recruit indicators is simply that the surveys were well designed for other purposes and provide insufficient coverage of the spatial and temporal scales of the juveniles (Albert et al., 2001; Ralph and Lipcius, 2014).

This paper focuses on the use of survey-based pre-recruit abundance indices and the degree of agreement between survey-based and stock assessment estimates of annual recruitment for species with juvenile coastal dependence. Accurate short-term forecasts of recruitment could improve the management advice in the stock assessment of species with juvenile coastal dependence. We focused on those ICES-assessed species whose juveniles rely on coastal habitats (see definitions in Seitz et al., 2014) and reviewed the use of survey-based pre-recruit abundance indices for short-term forecasts. For all ICES-assessed stocks whose juveniles use coastal habitats, we collated information from stock assessment reports and from a complementary questionnaire, which we designed for completion by the lead fisheries scientist for each stock assessment. The goals of our analysis were to: (1) assess the frequency

of the use of survey-based pre-recruit abundance indices in recruitment forecasts in the framework of ICES stock assessment working groups; (2) identify factors that influence when survey-based pre-recruit abundance indices are used; (3) determine the level of accuracy (agreement with stock assessment estimates) when survey-based pre-recruit abundance indices are used to indicate recruitment; and (4) suggest possible factors that influence the accuracy of the survey-based estimates. Our focus was on goals (1) and (3) because we had relatively high confidence in the underlying information and they provide important results about the frequency of use of pre-recruit surveys and their overall performance. The reliability of information to achieve goals (2) and (4) was uncertain, as it is difficult to judge a survey program for generating pre-recruit information when the survey was designed for other purposes (goal 2) and our sample size of surveys was too small for assessing which factors influence accuracy (goal 4).

Methods

Data collection

Of the 61 species for which ICES carried out stock assessments in 2017 and 2018, 18 species (Table 1) had juveniles with coastal dependence (Seitz et al., 2014). These 18 species encompass 78 distinct stocks. Information about the use of survey-based pre-recruit abundance indices for these ICES-assessed 78 stocks was collated. The information came from the ICES stock assessment working group (WG) reports (ICES, 2017a-c; ICES, 2018a-f), and the questionnaire completed by the lead fisheries scientists in charge of each stock assessment. The ICES WG reports, questionnaire responses, and follow-up communications with WG members provided the following information on the 78 stocks that rely on coastal habitat:

1. ICES DLS (data-limited stocks) category (ICES, 2012). The categories spanned from DLS category 1 (data-rich stocks with quantitative assessments) to DLS category 3 (stocks for

which survey-based assessments indicate trends) to categories DLS 4-6 (data-poor stocks without quantitative assessments).

2. Whether pre-recruit surveys were used for short-term estimation and prediction of recruitment. In ICES stock assessment WG terminology, recruitment estimation means projecting the youngest assessed year class strength for years y and $y+1$. The term recruitment prediction is used in WGs to calculate TAC advice when recruitment is projected two years ahead. In the present analysis, we pooled these two situations and considered the use of pre-recruit surveys both for recruitment estimation or prediction (hereafter called “short-term forecasts of recruitment”). Performing recruitment estimation is the minimum required and is mandatory for DLS category 1, but is highly unusual for the other categories.
3. Availability of survey-based abundance estimates for pre-recruits. The expertise of the lead fishery scientist involved with the assessment was the key source for these estimates. Indeed, WG reports only mention survey-based abundance indices when used in stock assessment. When they are not accounted for, expertise is the only means to investigate whether such indices exist.
4. When used, how were the short-term survey-based pre-recruit abundance indicators combined with the stock assessment? Survey-based pre-recruit abundance indices are typically used in two ways in ICES stock assessments: (i) post-hoc short-term forecasts of year-class strength by calibration-regression analysis of recruit index series (e.g., RCT3; Shepherd, 1997) and then used to account for future recruitment after a matrix model-based stock assessment is completed (e.g., Extended survivors analysis, XSA; Shepherd, 1999); or (ii) state-space modeling (e.g., SAM; Nielsen and Berg, 2014) that integrates the survey-based pre-recruit abundance indices directly into a stock assessment. We analyzed both uses of survey indices.

150 When survey-based pre-recruit abundance was available as an index (positive response to item
151 3 above), additional information was collated for that subset of stocks:

152 5. Sampling gear (i.e., acoustic, trawl or net) used in the survey to derive the pre-recruit index.

153 6. Spatial scale of the survey as one of four possibilities: (i) stock scale that included juvenile
154 habitats; (ii) stock scale that did not include juvenile habitats; (iii) stock spatial distribution
155 partially covered with the area covered including juvenile habitats; and (iv) stock distribution
156 partially covered and juvenile habitats not sampled.

157 7. Average number of samples in the annual survey.

158 8. Age-group represented in the survey-based recruitment estimate and the youngest age-group
159 included in the stock assessment.

160 Finally, when responses indicated that a stock assessment included short-term forecasts of
161 recruitment and a pre-recruit survey was available but not used to forecast recruitment:

162 9. The fisheries scientist for that stock assessment was asked why the survey was not used. Four
163 possible responses were offered in the questionnaire: (i) the pre-recruit index time series was
164 incomplete; (ii) the pre-recruit survey was carried out too late in the year to be available for
165 the ICES stock assessment working group; (iii) the potential use of the survey-based pre-
166 recruit abundance indices had not been evaluated; or (iv) pre-recruit survey-based indices
167 were investigated (e.g., during the benchmark procedure) but a decision was made to exclude
168 them from analysis.

169169

170 *Analysis: Availability and use of survey-based pre-recruit abundance indices for short-term*
171 *forecasting in assessment*

172 The frequency of the use of short-term forecasts of recruitment in stock assessment, and
173 the availability and the use of survey-based pre-recruit abundance indices to forecast
174 recruitment, were estimated from the WG reports and questionnaires collated for each stock.

Starting with the 78 (18 species) ICES-assessed stocks, we categorized these by habitat (demersal, benthic, pelagic). These stocks were further subdivided into those that used short-term forecasts in their assessments and either did or did not use available pre-recruit survey-based indices. For the subset of stocks that did not use the survey-based pre-recruit indices, the reasons for disuse by the WG assessors were noted. Another subset of stocks, that relied on short-term recruitment forecasts and also used pre-recruit survey results to generate short-term forecasts, was further analysed for accuracy of the survey-based predictions.

Analysis: Accuracy of survey-based pre-recruit abundance indices to forecast recruitment

Time series of survey-based recruitment predictions were obtained from ICES WG reports for each of the stocks that used survey-based pre-recruit indices for forecasting short-term recruitment in the assessment (ICES, 2017a-c; ICES, 2018a-f). For these stocks, time series of model-based recruitment short-term forecasts were obtained from the ICES database (ICES, 2018g). Complementary analyses were performed to assess the potential for autocorrelation between survey-based and model-based short-term forecasts of recruitment, because for some stocks the survey was also used within the assessment. When survey-based pre-recruit abundance indices were not used in the stock assessment modelling, but rather to make short-term forecasts post-assessment, the survey-based and stock-assessment-based indices were inherently independent and could be directly compared. However, when the survey-based pre-recruit abundance indices were used within the stock assessment, they influenced the assessment-based recruitment indices and could result in artificial agreement between the two short-term forecasts of recruitment because they were no longer independent.

Two alternative options were used to reduce or to remove this potential for artificial agreement between the two short-term forecasts (survey and assessment) of recruitment: (1) elimination of the last two years from the analysis, and (2) re-run of the stock assessment

without the survey index included to generate assessment-based recruitment not influenced by the survey results:

- The influence of survey results on assessment-generated estimates of recruitment can be significant, especially for the last years in a stock assessment (Hilborn and Walters, 1992). The influence of the survey results diminishes over time, as other sources of information in the stock assessment (e.g., catch-at-age and survey data on the older ages) inform the estimated recruitment values. To partially account for dependence between the survey- and model-based estimates, we eliminated the last two years of recruitment estimates for those stocks that used the survey-derived estimates as part of their stock assessment modeling. This elimination was done either manually or because the last two years were dropped when matching the two recruitment indices (i.e., there were no survey estimates available to match recruitment for the last two years of the assessment). To test the robustness of these modelling option, we employed two methods, both of which focus on the accuracy of the correlation results from stocks that used survey indices in their assessments: The first was a comparison between the four stocks with independent survey and assessment estimates of recruitment and the remaining 10 stocks that included the survey index in their assessment. The second was a windowing approach to compute correlations between survey and assessment estimates of recruitment, to assess the influence of the last years in correlations (see details in supp. Mat. 2).

- The best way to address this potential for artificial agreement is to re-run the stock assessments without the survey-derived indices, and then compare the new assessment-based estimated recruitments with the, now independent, survey-derived estimates of recruitment. Such an approach is obviously the most attractive in theory, but each assessment varies among the different stocks and cannot be tuned from the ICES database without the expertise of the stock assessment WG. To do so, the fisheries scientists in charge of these stock assessments were asked to re-run the stock assessments without the survey-derived indices, and some of

225 them kindly did so. These new time series of model-based recruitment were collated and used
226 separately from the potentially correlated estimates in analyses. This subset of comparisons
227 allowed us to evaluate the robustness of results based on the potentially correlated estimates.
228 For standardization purpose, we also eliminated the last two years of the recruitment estimates
229 from these series, either manually or naturally.

230 To assess the accuracy of the survey-based predictions of recruitment compared to
231 assessment-based estimates, we computed the Pearson correlation coefficient (r) between the
232 survey-based recruitment estimates and the stock assessment model-based abundance for the
233 youngest year group. This was done for all stocks (r_1 , using model-based data from ICES
234 database) and for the subset of 10 stocks that the assessment estimates were independent of the
235 survey (r_2 , from stocks whose assessment did not use survey or from re-run assessment models).
236 We assumed that the model-based estimates were a realistic value and thus the closer the
237 correlation of the survey-based prediction to the model-based value, the higher the accuracy of
238 the survey-based value. Because the true value of recruitment is unknown, we refer to this as
239 apparent accuracy. While agreement between the two estimates of recruitment suggest higher
240 confidence in the survey-based estimates, without knowing the true values of recruitment we
241 cannot access whether either is or both are biased.

242 For the stocks for which correlation coefficient r_1 (model-based data from ICES
243 database) and r_2 (for rerun assessment estimates) were available, we first compared their
244 respective levels to highlight potential lack of independence and caution about interpretation of
245 r_1 . From this preliminary analysis (r_1 versus r_2 for rerun stocks only), we determined if we would
246 use the r_2 values (truly independent estimates on 10 stocks) rather than the r_1 (14 stocks but
247 only 4 trully independent estimates only) in subsequent analyses.

248 Another proxy (r_3) was designed to approximate how short-term recruitment forecasts
249 can be used in stock assessments that do not have a source of year-specific short-term forecasts.

The geometric mean of the model-based abundances for the youngest year class during the previous five years was computed. When year-specific forecasts of recruitment are not used, geometric mean of model-based recruitment estimates is frequently used in forecasting for ICES stock assessments. To estimate the improvement of the forecast linked to the use of survey-based pre-recruit abundance indices, r_1 or r_2 and r_3 were compared. We used a one-way analysis of variance (ANOVA), after an arcsine transformation, to compare r_1 or r_2 to r_3 values. The arcsine transformation is appropriate to normalize the data from the original $[-1,1]$ distribution of correlation coefficients (Sokal and Rohlf, 1995). A higher value of r_1 or r_2 (for the survey-based estimates) compared to r_3 (geometric mean of the assessment-based estimates) indicates that survey estimates agree with assessment values better than average recruitment agrees with the assessment values. In this way, r_3 is an approximate proxy of the contribution of survey-based pre-recruit indices to estimate future recruitment over and above the use of a 5-year average.

We explored whether various factors influenced the magnitude of r_1 or r_2 , including species vertical guild (Table 1), sampling gear, scale of the survey, number of samples in the survey, age group in the survey-based pre-recruit abundance indices, youngest age group in the stock assessment, difference between these two ages, and length of the time series.

Results

Stocks of coastal dependent species

ICES performed stock assessments for 185 stocks in 2017-2018 that spanned 61 species. Eighteen of these species (30%), which involved 78 stocks (42%), depend on coastal juvenile habitat (Table 2; supp. Table 1). These 78 stocks are widespread in the North East Atlantic (from Iberian waters to Greenland in latitude and from the North Sea to Greenland in longitude) and in the Baltic Sea (supp. Table 1). The habitat use of these species and stocks with juvenile

coastal dependence were: demersal (9 species; 39 stocks), benthic (6 species; 23 stocks), and pelagic (3 species; 16 stocks). Among these 78 stocks, most (87%) were well-assessed stocks (ICES categories 1 and 3), whereas 10% were data-poor stocks, all of which were demersal species (supp. Table 1).

Use of recruitment forecasts and pre-recruit surveys in assessment

Among the 78 stocks from species with juvenile coastal dependence, 49 (Table 2) used short-term recruitment forecasts (from any source) in their assessments. Most of these 49 stocks (46) were designated as DLS Category 1, with the remaining three stocks being DLS 3. Survey-based pre-recruit abundance indices were available (used and not used in the assessment) for 35 (71%) of these 49 stocks, which were all designated as DLS Category 1 (Table 2; Figure 1). For these 35 (of 78) stocks with both survey-based pre-recruit abundance indices available and that use short-term recruitment forecasts in their assessment (Table 2), the pre-recruit indices were derived mainly (supp. Table 1) from trawl surveys for demersal species (12 of 18 stocks) and benthic species (9 of 9 stocks), and from acoustic surveys for pelagic species (5 of 8 stocks). While survey-based pre-recruit abundance indices were available for 35 of the 49 stocks that generated recruitment forecasts in their assessments, only 14 of these 35 stocks (40%; Table 2; Figure 1) actually used the indices in their assessments. For the majority of stocks (21 of 35), the indices were not used for short-term forecasts of recruitment. The underutilisation of survey-based indices was noteworthy for stocks of demersal species (12 of 18 stocks did not use the indices; supp. Table 1).

Six stocks with unused indices reported that the available time series were not yet sufficient or because the results would not be available in time for consideration by the WG (Table 3). But, the most commonly reported reason for not using the survey-based indices (11 of 21) was that the use of the indices had not been thoroughly evaluated (Table 3; supp. Table

1). The remaining four stocks with unused indices had attempted to use the indices but a decision was made to not use them because the surveys were not designed to estimate pre-recruit abundance in the spatial domain of the stock (Table 3). A partial explanation for not using the survey-based indices when they were sufficient and available (15/21) was that these surveys were not designed to cover both the spatial scale of the stock and/or coastal juvenile habitats (Table 3).

Fourteen stocks used the survey-based pre-recruit indices in their forecasts. These 14 stocks are distributed in the North East Atlantic (from Bay of Biscay to Greenland in latitude and from the North Sea to Greenland in longitude) and in the Baltic Sea (Table 4). For these 14 stocks, seven of the indices were derived from surveys covering both the stock scale and coastal nurseries, four indices were from surveys that partially cover the stock's spatial extent and include coastal nurseries, and three indices were calculated from surveys done at the stock spatial scale but which do not include coastal juvenile habitat (Table 4).

Apparent accuracy of survey-based pre-recruit indices

For 12 of 14 stocks (Table 4), pre-recruit abundance indices were used in the assessments. These were either derived from a single survey (8 stocks,) or were combined into a single recruitment index as part of the assessment by the ICES working group (4 stocks, North Sea cod and sole, Irish Sea plaice and Celtic Sea whiting; ICES, 2017c). Two (of 14) stocks used two survey-based pre-recruit abundance indices for short-term forecasting (Table 4): Iceland cod (ICES, 2018c) and North Sea whiting (ICES, 2017c). Our analysis of the relationship between the survey-based pre-recruit abundance indices and the model-based abundance for the youngest year class (r_1 and r_2) considered a single survey-based pre-recruit abundance index of recruitment per stock. For North Sea whiting, the lead fishery scientist (Miethe, pers. com.) for the stock assessment (ICES, 2017c) indicated that the index in Autumn

(IBTSQ3) is considered as the reference pre-recruit abundance index. For Iceland cod, we initially analysed both indices separately (surveys SMB and SMH had correlation coefficients of 0.75 and 0.8 with model-based indices, respectively); given the similarity of the results, the SMH index derived from the fall survey was selected (Table 4).

Among these 14 stocks, four use survey-based pre-recruit abundance indices only for forecasting and 10 used these indices in both stock assessment and forecasting (Table 4). Of these 10, only five required manual deleting of two recent years. The other five stocks, which used the survey indices in their assessments, had sufficient lag between the age of fish in the survey and the age of recruitment (youngest age) in the assessment. This meant that the two most recent years of recruitment from the stock assessment would not be auto-correlated with their survey index for our comparisons (i.e. “Natural removal”, Table 4).

From the 10 stocks utilising survey-based indices in both stock assessment and forecasting, fisheries scientists in charge of assessments agreed to rerun the stock assessments without the survey-derived indices for six stocks (Table 4, r_2 in bold). For these stocks, correlations were higher for r_1 than for r_2 (Table 4, for the 6 stocks, average difference in Pearson correlation coefficient $r_1 - r_2 = 0.077$ [0, 0.19]). These patterns confirmed the preliminary tests of robustness on the use of the correlation between the survey-based recruitment estimates and the stock assessment model-based abundance; i.e., low to moderate influence of autocorrelation when the last two years of the recruitment estimates are removed (detailed in supp. Mat. 2). However slight, these differences did indicate an overestimation of r_1 through correlation induced by inclusion in the assessment. Hence, we selected r_2 for further analyses, which reduced the number of stocks to 10 (4 whose assessment did not use the index and 6 rerun assessments, Table 4).

When used, the survey-based predictions of recruitment (r_2) had a reasonable apparent accuracy (Table 4; Figure 2; Figure 3). Survey-based pre-recruit abundance indices had

significantly higher correlations with the model-based recruitment estimates than the geometric means of the five previous years of model-based abundances (Figure 3; $p < 0.001$, after arcsine transformations of r_2 and r_3). No obvious patterns emerged from the factors (species habitat, survey design, Table 4) that could influence the accuracy of the survey-based pre-recruit abundance indices r_2 , although the small size of the data set and the many potential influential factors made identification of associations difficult.

Discussion

We examined ICES-assessed stocks that both utilize coastal areas as juvenile habitat and use survey-based predictions of recruitment in their management assessments. Of the 78 stocks involving 18 species with juvenile coastal-dependence, 49 also used short-term forecasts of recruitment in assessments. Most of these stocks (46 of 49) were designated as ICES DLS Category 1 stocks. Indeed, short-term forecasts of recruitment are mandatory in the ICES protocol for this category. We analysed the existence and aspects of surveys and derived survey-based pre-recruit indices and how they are presently used in assessments for the 78 stocks, using data collated from WG reports, responses to a questionnaire from the lead fishery scientists for each stock, and communications with lead members of various stock assessment WGs. We sought to explore how surveys are used to generate recruitment indices as part of assessments, possible reasons for their omission, and the accuracy of predicted recruitment from survey-derived values.

The responses to the questionnaire as to why the survey information was available but not used (i.e., survey data on pre-recruit abundance were not used for 21/35 = 60% of the stocks for which they are available) indicated that there are opportunities for determination of how the survey information, either as is or with some adjustments to the survey design, could be used in assessments. The most common response for why an available survey was not used was that

its utility had not been rigorously evaluated, followed by issues of whether enough data were available and that the survey results were not available in time for assessments. These three reasons accounted for why 17 of 21 stocks were not using available surveys to forecast recruitment for assessment, and suggest that surveys are available that, with proper evaluation, may be useful for generating recruitment indices.

Fishery-independent surveys are designed to answer specific questions and their lack of use for other purposes is not indicative of a poorly designed survey. For our proposed use, to forecast recruitment, the coverage of coastal habitats and the effective sampling of pre-recruit juveniles are critical. Both the stocks that did not use surveys to predict recruitment and those that did confirmed the (perhaps obvious) importance of the spatial scales of the surveys. Half of the survey-based pre-recruit indices used in assessments covered both the stock scale and coastal juvenile habitat, while the other half covered either stock scale or juvenile habitats. In contrast, none of the unused survey-based pre-recruit abundance indices covered both the stock scale and the coastal juvenile habitat. Most (87%) of the unused pre-recruit abundance survey-based indices covered only a fraction of the spatial extent of the stock, and 47% did not sample coastal juvenile habitat.

A major challenge for estimating pre-recruit abundance indices from surveys is to account for complex spatial and temporal variation in pre-recruit abundance (Denson et al., 2017; Potts and Rose, 2018). Variation in abundance across successive juvenile stages could be driven by small scale processes, leading to large spatial discrepancies among juvenile habitats (Scharf, 2000). The temporal (including inter-annual) variability in coastal habitat use of juvenile fish suggests that to estimate recruitment, it is necessary to survey several juvenile habitats (Chittaro et al., 2008). Both juvenile coastal distributions outside the geographical area covered by the surveys and regional patterns in recruitment variability (Denson et al., 2017)

may hinder estimation of reliable recruitment estimates (Albert et al., 2001; Ralph and Lipcius, 2014).

The 17 stocks with available surveys not being used and which have not been evaluated for use would need to be evaluated. The evaluation should consider whether the sampling design can generate sufficiently accurate predictions of recruitment, and how easy it would be to maintain present sampling and make minor additions to better cover nursery areas (e.g., add stations in shallow juvenile habitat). Thus, there is an opportunity for further analyses to determine the feasibility and utility of these surveys for also generating short-term forecasts of recruitment, either as they are presently implemented or with minor changes that do not affect the use of the surveys for other purposes.

When survey-based predictions of recruitment were used in assessments, their apparent accuracy was reasonably high. The r_2 values averaged 0.76 across all 10 stocks. Such degree of agreement was based on stocks with independent survey and assessment estimates and therefore was not influenced by lack of independence due to use of surveys within assessments. Indeed, for four stocks, survey-based predictions of recruitment were originally independent of the assessments (Table 4). For the six remaining stocks, models were rerun after removing survey-based indices from the assessment. For these six stocks differences between r_1 and r_2 depended at least partly on the availability of alternative information on recruitment strength used in stock assessment models. The difference was insignificant for North Sea plaice, for which several alternative data-based sources of information are used in the assessment model to infer pre-recruit abundance (including survey-based indices from other surveys; ICES, 2017c). Conversely, r_1 - r_2 reached 0.19 for the western Baltic Sea cod, for which recruitment is mainly informed by the survey-based index in the assessment model for young stages (ICES, 2018b). This difference illustrates autocorrelation between survey-based and model-based short-term

forecasts of recruitment; i.e., for stocks where the survey-based recruitment indices informed the assessment models.

The degree of agreement between survey-based and survey-independent, model-based short-term forecasts was not due to a few influential points, as there was an average of 22 years in the various time series. Furthermore, the survey-based predictions out-performed the alternative using a 5-year geometric mean of model-based values.

Given the long history of attempts to predict recruitment in fisheries management, our results strongly suggest that juvenile surveys should be investigated for their potential use in assessments; a theme that has been emphasized by analysis of other stocks (Helle et al., 2000; Zhang et al., 2010; Caputi et al., 2014; Punt, 2019). Any possible use of survey results would need to be evaluated for the specifics of the survey data, the assessment methodology, and the life history of the species.

Deviations between survey-based and model-based short-term forecasts of recruitment may be due to several factors. First is the unknown estimation error in deriving recruitment estimates from surveys due to high spatio-temporal variation in abundance (Denson et al., 2017; Potts and Rose, 2018). Quantifying and understanding the causes of these errors is central to obtaining reliable recruitment estimates (Albert et al., 2001; Ralph and Lipcius, 2014). Second, our assumption that the model-based estimates are accurate ignores how process and estimation errors in recruitment arise from stock assessment models (Hilborn and Walters, 1992). Estimates of recruitment time-series are sensitive to model assumptions used in the assessments (Dickey-Collas et al., 2015). Third, there may be high, density-dependent and variable juvenile mortality (Nash et al., 2007; Le Pape and Bonhommeau, 2015; Haggarty et al., 2017) after the survey-based estimate of pre-recruit abundance. Given that these and other factors add noise to both survey-based and model-based short-term forecasts of recruitment, the degree of

agreement we found between both predictors across diverse stocks and sampling programs is encouraging.

The small (10 stocks) dataset precluded a comprehensive analysis of the driving factors of survey apparent accuracy. The correlation values did not indicate any obvious dependence on species habitat nor survey design. However, these and other factors, such as life history of the species, probably influence survey accuracy, which warrants analysis with more stocks. Two main issues complicated our ability to determine the factors that influenced the accuracy of survey-based pre-recruit estimates: (i) it is speculative to judge a survey program for generating pre-recruit information when the survey was designed for other purposes, and (ii) our sample size was too small for using the questionnaire results for assessing which factors influence accuracy. Given these caveats, the present analysis allows for some recommendations about survey design to ensure that the surveys provide sufficiently accurate pre-recruit abundance indices for advice about recruitment in stock assessment of species with juvenile coastal dependence:

- Surveys should sample coastal juvenile areas at appropriate times, to avoid the high and variable mortality during the early juvenile stages (Nash et al., 2007; Le Pape and Bonhommeau, 2015; Haggarty et al., 2017).
- Surveys should cover a large proportion of a stock's spatial domain to capture inter-annual variation in nursery habitat utilization (Albert et al., 2001; Ralph and Lipcius, 2014).
- Surveys should be carried out annually to avoid missing values in the pre-recruit abundance time series.
- The juvenile portion of the survey should include an evaluation of the performance of the sampling gear (e.g., selectivities) and incorporate methods for quantifying variability.

- Where possible, juvenile surveys or the juvenile component of stock surveys should aim to be as consistent as possible with the survey of non-juvenile areas to provide commensurable data for combined analyses.

These conditions provide a general basis for examining how surveys can be initially evaluated for possible use for juveniles and pre-recruit indices. These recommendations can be applied to situations when surveys are being revised (surveys are presently done for multiple reasons) and new surveys are being designed.

Augmenting the survey-based pre-recruit abundance indices with other covariate variables, such as environmental drivers, may further improve the accuracy of recruitment predictions. Indices based on environmental drivers (e.g., ICES, 2018a for North East Arctic cod; Le Pape et al., 2003 and Lagarde et al., 2018 for Bay of Biscay sole; Denson et al., 2017) alone, or in combination with pre-recruit abundance indices (Zhang et al., 2010; Ralston et al., 2013), could provide helpful information about recruitment trends and variability in the near term. However, changes in total allowable catch (TAC) recommendations lead to gains only when environmental predictors and survey-based pre-recruit abundance indices are accurately assessed (Basson, 1999; De Oliveira and Butterworth, 2005). The increase in accuracy that survey-based pre-recruit abundance indices can provide to catch advice suggests that existing surveys should be evaluated for their potential use.

Predictions of future short-term recruitment can influence management advice both for the assessment year and for the TAC year (ICES, 2015). Our analysis showed that, while a limited number of the total possible stocks that can use survey-based predictions actually use them, when survey-based predictions are used in the assessment their apparent accuracy is reasonable. Survey-based pre-recruit abundance indices are being used for some stocks either explicitly in the stock assessment model (e.g., SAM model; Nielsen and Berg, 2014), or in a separate forecasting routine combined with stock assessment outputs (e.g., RCT3 routine post

XSA model; Shepherd 1997; Shepherd, 1999). These indices inform the expected recruitment in future years. The scope of the present paper was focused on the usefulness of survey-based pre-recruit abundance indices for advice about recruitment, not on the ways in which to utilise these indices in stock assessment procedures; this has been extensively discussed by others (Punt, 2019).

Tools for forecasting recruitment play an important role in fisheries management and decision-making, and all possible tools should be at least explored for their potential utility, if not utilised. When catches are highly dependent on recruitment (short-lived or over-exploited stocks; e.g., North Sea cod, ICES, 2017c), estimating recruitment and possible variability about the forecast is a priority to provide reliable information for management. However, the number of years for which short-term forecasts can benefit from survey-based abundance indices of pre-recruits obviously depend on the year-lag between the first age in the catch forecast and the age of the pre-recruit individuals in the survey. For the large proportion of stocks with only a 1-year lag (Supp. Table 1), there is no observed recruitment survey index for more years ahead, and short-term forecast means a forecast for the next year only.

Even when they are not accounted for in stock assessment, survey-based pre-recruit abundance indices could be considered as quantitative evidence supporting or opposing predictions derived using average previous recruitment, and used to provide a measure of the uncertainty in predicted recruitment. Indeed, when the survey-based pre-recruit abundance indices are not available during an assessment (e.g., Sandeel stocks, sup. Table 1; Table 3), some procedures allow their results to be considered *a posteriori*. For example, the advice for the main flatfish and round fish stocks in the North Sea has a procedure for reopening after the surveys are conducted in autumn (ICES, 2008; ICES, 2015). Re-evaluating management advice after surveys are completed and pre-recruit abundance indices are estimated to differ significantly from assessment derived indices should make the advice more robust (ICES,

2008). This procedure of re-evaluating management advice clearly shows the validity and importance of the recruitment indices. We recognize that these approaches introduce additional work for those delivering advice; thus, exploratory analyses to assess their potential benefits to assessments are a good first step. While our focus was on species that use coastal habitats, our evaluation approach is applicable to most species, including those that do not depend on coastal juvenile habitats (Kimoto et al., 2007; Ralston et al., 2013).

We focused our analysis on using existing surveys for stocks that use recruitment forecasts in their assessments. In addition to the use of survey-based pre-recruit abundance indices for forecasting recruitment, fishery-independent surveys can be evaluated for their potential use with other management goals. Examples include quantifying juvenile habitat for informing an ecosystem-based approach to fisheries management (Browman et al., 2004), deriving indices of environmental drivers for further forecasting (Hidalgo et al., 2016), and for informing dynamic marine spatial plans that respond to changes in coastal habitats (Kininmonth et al., 2019). Surveys can also be used to provide alerts on the impacts of anthropogenic disturbances affecting survival of juveniles. A large proportion of coastal-dependent species is impacted by human activity other than fishing mortality when juveniles utilize coastal habitats (Brown et al., 2018a). Regular monitoring of juvenile habitats to provide data for assessment can generate spatially-explicit evidence for local productive areas to inform environmental management. Surveys can provide information on juvenile responses to both environmental drivers (Hermant et al., 2010; Caputi et al., 2014; Lagarde et al., 2018; Brown et al., 2019) and anthropogenic pressures (Rochette et al., 2010; Archambault et al., 2018), which can influence future stock dynamics (Stige et al., 2013). Habitat degradation can result in either overly optimistic or overly conservative assessments of stock status (Brown et al., 2018b). Preserving or restoring the capacity of juvenile habitat is of major importance for improving adult biomass of populations relying on coastal juvenile habitat (Van de Wolfshaar et al., 2011; Le Pape and

Bonhommeau, 2015; Archambault et al., 2018). Existing and planned surveys should be examined for possible leveraging of their results, in addition to their primary motivation and goals, thereby integrating fisheries and ecosystem-based management (Kraufvelin et al., 2018).

550550

551 Acknowledgements

This work was developed within the context of the ICES working group WGVHES (Working Group on the Value of Coastal Habitats for Exploited Species). The authors thank both ICES and all participants of the working group 2017-2019. The authors also warmly thank Maria Lifentseva and Jette Fredslund (ICES) for their efficient help to connect us to scientists in charge of the 78 stock assessments. The authors thank Mark Dickey-Collas (ICES) and the scientists involved in stock assessments for their contributions. Special thanks to Marianne Robert, Niels Hintzen, and Marie Storr-Paulsen who kindly and greatly contributed by re-tuning stock assessment models without the recruitment index data. Finally, the authors thank Dr. Stan Kotwicki, the editor, Niels Hintzen and the 2 other anonymous reviewers for their constructive reviews that improved the manuscript.

562562

Supplementary Material is available at ICES JMS online

564564

565565

566 References

567567

Albert, O.T., Nilssen, E.M., Nedreaas, K.H. and Gundersen, A.C. 2001. Distribution and abundance of juvenile North-East Arctic Greenland halibut (*Reinhardtius hippoglossoides*) in relation to survey coverage in the physical environment. ICES Journal of Marine Science 58: 1053-1062.

572

573 Archambault, B., Le Pape, O., Bousquet, N. and Rivot, E. 2014. Density dependence can be
574 revealed by modeling the variance in the stock-recruitment process. An application to flatfishes.
575 ICES Journal of Marine Science 71: 2127-2140.

576

577 Archambault, B., Rivot, E., Savina, M. and Le Pape, O. 2018. Using a spatially structured life
578 cycle model to assess the influence of multiple stressors on an exploited coastal-nursery-
579 dependent population. Estuarine Coastal and Shelf Science 201: 95-104.

580

581 Basson, M. 1999. The importance of environmental factors in the design of management
582 procedures. ICES Journal of Marine Science 56: 933-942.

583

584 Beck, M.W., Heck, K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M.,
585 Halpern, B., Hays, C.G., Hostino, K., Minello, T.J., Orth, R.J., Sheridan, P., and Weinstein,
586 M.P. 2001. The identification, conservation and management of estuarine and marine nurseries
587 for fish and invertebrates. Bioscience 51: 633-641.

588

589 Brown, E.J., Vasconcelos, R.P., Wennhage, H., Bergström, U., Støttrup, J.G., van de
590 Wolfshaar; K., Millisenda, G., Colloca, F. and Le Pape, O. 2018a. Conflicts in the coastal zone:
591 A rapid assessment of human impacts on commercially important fish species utilizing coastal
592 habitat. ICES Journal of Marine Science 75: 1203–1213.

593

594 Brown, E.J., Kokkalis, A., Støttrup, J.G. 2019. Juvenile fish habitat across the inner Danish
595 waters: Habitat association models and habitat growth models for European plaice, flounder
596 and common sole informed by a targeted survey. Journal of Sea Research 155: 1-16.

597

598 Brown, J.C., Broadley, A., Adame, M.F., Branch, T., Turcschwell, M.P. and Connolly, R.M.
599 2018b, in press. The assessment of fishery status on fish habitats. *Fish and Fisheries* 20: 1-14.

600

601 Browman, H.I., Stergiou, K.I., Cury, P.M., Hilborn, R., Jennings, S., Lotze, H.K., Mace, P.M.,
602 Murawski, S., Pauly, D., Sissenwine, M. and Zeller, D. 2004. Perspectives on ecosystem-based
603 approaches to the management of marine resources. *Marine Ecology Progress Series* 274: 269–
604 303.

605

606 Caputi, N., de Lestang, S., Hart, A., Kangas, K., Johnston, D., Penn, J. 2014. Catch predictions
607 instock assessment and management of invertebrates fisheries using pre-recruit abundance -
608 Case studies from western Australia. *Reviews in Fisheries Science & Aquaculture* 22: 36-54.

609

610 Cowan, J.H., Rose, K.A. and de Vries, D.R. 2000. Is density dependent growth in young of the
611 year fishes a question of critical weight? *Reviews in Fish Biology and Fisheries* 10: 61-89.

612

613 Cury, P.M., Fromentin, J.M., Figuet, S. and Bonhommeau, S., 2014. Resolving Hjort's
614 dilemma: How is recruitment related to spawning stock biomass in marine fish? *Oceanography*
615 27: 42-47.

616

617 De Oliveira, J.A.A. and Butterworth, D.S., 2005. Limits to the use of environmental indices to
618 reduce risk and/or increase yield in the South African anchovy fishery. *African Journal of*
619 *Marine Science* 27: 191-203.

620

621 Denson, L.S., Sampson, D.B. and Stephens, A. 2017. Data needs and spatial structure
622 considerations in stock assessments with regional differences in recruitment and exploitation.
623 Canadian Journal of Fisheries and Aquatic Sciences 74: 1918-1929.

624

625 Dickey-Collas, M., Hintzen, N.T., Nash, R.D., Schon, P.J., Payne, M.R. 2015. Quirky
626 patterns in time-series of estimates of recruitment could be artefacts. ICES Journal of Marine
627 Science 72: 111-116.

628

629 Dingsor, G.E., Cianelli, L., Chan, K.S., Ottersen, G. and Stenset, N.C. 2007. Density
630 dependence and density independence during the early life stages of four marine fish stocks.
631 Ecology 88: 625-634.

632

633 Haggarty, D.R., Lotterhos, K.E. and Shurin, J.B. 2017. Young-of-the-year recruitment does not
634 predict the abundance of older age classes in black rockfish in Barkley Sound, British
635 Columbia, Canada. Marine Ecology Progress Series 574: 113-126.

636

637 Helle, K., Bogstad, B., Marshall, C.T., Michalsen, K., Ottersen, G. and Pennington, M., 2000.
638 An evaluation of recruitment indices for Arcto-Norwegian cod (*Gadus morhua* L.). Fisheries
639 Research 48: 55-67.

640

641 Hermant, M., Lobry, J., Poulard, J.C., Désaunay, Y., Bonhommeau, S. and Le Pape O. 2010.
642 Impact of warming on abundance and occurrence of flatfish populations in the Bay of Biscay
643 (France). Journal of Sea Research 64: 45-53.

644

645 Hilborn, R. and Walters, C. 1992. Quantitative Fisheries Stock Assessment: Choice, Dynamics
646 and Uncertainty. Springer Nature. 570 pp.

647

648 Hidalgo, M., Secor, D.H. and Browman, H.I. 2016. Observing and managing seascapes: linking
649 synoptic oceanography, ecological processes, and geospatial modelling. ICES Journal of
650 Marine Science 73: 1825–1830.

651

652 Houde, E.D. 2008. Emerging from Hjort's shadow. Journal of the Northwest Atlantic Fisheries
653 Society 41: 53-70.

654

655 ICES. 2008. Report of the Ad hoc Group on Criteria for Reopening Fisheries Advice
656 (AGCREFA). ICES CM 2008/ACOM: 60.

657

658 ICES. 2012. DLS Guidance Report, ICES Implementation of Advice for Data-limited Stocks
659 in 2012 in its 2012 Advice. ICES Advisory committee. ICES CM 2012/ACOM: 68, 42 pp.

660

661 ICES. 2015. Report of the Benchmark Workshop on North Sea Stocks (WKNSEA), 2–6
662 February 2015, Copenhagen, Denmark. ICES CM 2015/ACOM: 32, 253 pp.

663

664 ICES. 2017a. Report of the Working Group on Widely Distributed Stocks (WGWIDE), 30
665 August–5 September 2017, ICES Headquarters, Copenhagen, Denmark. ICES CM
666 2017/ACOM: 23, 1111 pp.

667

668 ICES. 2017b. Report of the Working Group on Celtic Seas Ecoregion (WGCSE), 9–18 May
669 2017, Copenhagen, Denmark. ICES CM 2017/ACOM: 13, 1464 pp.

670

671 ICES. 2017c. Report of the Working Group on Assessment of Demersal Stocks in the North
672 Sea and Skagerrak (NSSK), 26 April–5 May 2017, ICES HQ. ICES CM 2017/ACOM: 21, 1248
673 pp.

674

675 ICES. 2018a. Report of the Arctic Fisheries Working Group (AFWG), 18–24 April 2018, Ispra,
676 Italy. ICES CM 2018/ACOM: 06, 857 pp.

677

678 ICES. 2018b. Report of the Herring Assessment Working Group for the Area South of 62°N
679 (HAWG). 29–31 January 2018 and 12–20 March 2018. ICES HQ, Copenhagen, Denmark.
680 ICES CM 2018/ACOM: 07, 958 pp.

681

682 ICES. 2018c. Report of the North-Western Working Group (NWWG), 26 April–3 May 2018,
683 ICES HQ, Copenhagen, Denmark. ICES CM 2018/ACOM: 09, 733 pp.

684

685 ICES. 2018d. Baltic Fisheries Assessment Working Group (WGBFAS), 6–13 April 2018, ICES
686 HQ, Copenhagen, Denmark. 727 pp.

687

688 ICES. 2018e. Report of the Working Group for the Bay of Biscay and the Iberian Waters
689 Ecoregion (WGBIE), 3–10 May 2018, ICES HQ, Copenhagen, Denmark. ICES CM
690 2018/ACOM: 12, 642 pp.

691

692 ICES. 2018f. Report of the Working Group on Southern Horse Mackerel, Anchovy and Sardine
693 (WGHANSA) 26–30 June 2018, Lisbon, Portugal. ICES CM 2018/ACOM: 17, 597 pp.

694

- 695 ICES. 2018g. ICES Stock Assessment Database. Copenhagen, Denmark. ICES. 2018/01/01.
696 <http://standardgraphs.ices.dk>.
697
- 698 Juanes, F. 2007. Role of habitat in mediating mortality during the post-settlement transition
699 phase of temperate marine fishes. *Journal of Fish Biology* 70: 661-677.
700
- 701 Kimoto, A., Mouri, T. and Matsuishi, T. 2007. Modelling stock–recruitment relationships to
702 examine stock management policies. *ICES Journal of Marine Science* 64: 870–877.
703
- 704 Kininmonth, S., Weeks, R., Abesamis, R.A., Bernardo, L.P.C., Beger, M., Treml, E.A.,
705 Williamson, D. and Pressey, R.L. 2019. Strategies in scheduling marine protected area
706 establishment in a network system. *Ecological Applications* 29: 1-10.
707
- 708 Kraufvelin, P., Pekcan-Hekim, Z., Bergstrom, U., Florin, A.B., Lehikoinen, A., Mattila, J.,
709 Arula, T., Briekmane, L., Brown, E.J., Celmer, Z., Dainys, J., Jokinen, H., Kaaria, P.,
710 Kallasvu, M., Lappalainen, A., Lozys, L., Moller, P., Orio, A., Rohtla, M., Saks, L., Snickars,
711 M., Støttrup, G., Sundblad, G., Taal, I., Ustups, D., Verliin, A., Vetemaa, M., Winkler, H.,
712 Wozniczka, A. and Olsson, J. 2018. Essential coastal habitats for fish in the Baltic Sea.
713 *Estuarine, Coastal and Shelf Science* 204: 14-30.
714
- 715 Lagarde, A., Doyen, L., Ahad-Cissé, A., Gourguet, S., Le Pape, O., Thébaud, O., Caill-Milly,
716 N., Morandeau, G. and Macher, C. 2018. How does MMEY mitigate the bioeconomic effects
717 of climate change for mixed fisheries. *Ecological Economics* 154: 317-332.
718

- 719 Le Pape, O., Chauvet, F., Mahévas, S., Lazure, L., Guérault, G. and Désaunay, Y. 2003.
720 Quantitative description of habitat suitability for the juvenile common sole (*Solea solea*, L.)
721 and contribution of different habitats to the adult population in the Bay of Biscay (France).
722 Journal of Sea Research 50: 139-149.
- 723
- 724 Le Pape, O. and Bonhommeau, S. 2015. The food limitation hypothesis for juvenile marine
725 fish. Fish and Fisheries 16: 373-398.
- 726
- 727 Levin, P.S. and Stunz, G.W. 2005. Habitat triage for exploited fishes: can we identify essential
728 fish habitat? Estuarine, Coastal and Shelf Science 64: 70-78.
- 729
- 730 Nash, R.D.M., Geffen, A.J., Burrows, M.T. and Gibson, R.N. 2007. Dynamics of shallow-water
731 juvenile flatfish nursery grounds: application of the shelf-thinning rule. Marine Ecology
732 Progress Series 344: 231-244.
- 733
- 734 Lorenzen, K., and Camp, E.V. 2019. Density-dependence in the life history of fishes: when is
735 a fish recruited? Fisheries Research 217: 5-10.
- 736
- 737 Needle, C.L. 2001. Recruitment models: diagnosis and prognosis. Reviews in Fish Biology and
738 Fisheries 11: 95-111.
- 739
- 740 Nielsen, M. and Berg, C. 2014. Estimation of time-varying selectivity in stock assessments
741 using state-space models. Fisheries Research 158: 96-101.
- 742
- 743 Potts, S.E. and Rose, K.A. 2018. Evaluation of GLM and GAM for estimating population

744 indices from fishery independent surveys. Fisheries Research 208: 167-178.

745

746 Punt, A.E. 2019. Recruitment: theory, estimation, and application in fishery stock assessment
747 models. Fisheries Research 217: 1-4.

748

749 Ralph, G.M. and Lipcius, R.N. 2014. Critical habitats and stock assessment: age-specific bias
750 in the Chesapeake Bay blue crab population survey. Transactions of the American Fisheries
751 Society 143: 889-898.

752

753 Ralston, S., Sakuma, K.M. and Field, J.C., 2013. Interannual variation in pelagic juvenile
754 rockfish (*Sebastes* spp.) abundance—going with the flow. Fisheries Oceanography 22: 288-308.

755

756 Rochette, S., Rivot, E., Morin, J., Mackinson, S., Riou, P. and Le Pape O. 2010. Effect of
757 nursery habitat destruction on flatfish population renewal. Application to common sole (*Solea*
758 *solea*, L.) in the Eastern Channel (Western Europe). Journal of Sea Research 64: 34-44.

759

760 Seitz, R.D., Wennhage, H., Bergstrom, U., Lipcius, R.N. and Ysebaert, T. 2014. Ecological
761 value of coastal habitats for commercially and ecologically important species. ICES Journal of
762 Marine Science 71: 648-655.

763

764 Scharf, F. 2000. Patterns in abundance, growth, and mortality of juvenile red drum across
765 estuaries on the Texas coast with implications for recruitment and stock enhancement.
766 Transactions of the American Fisheries Society 129: 1207-1222.

767

768 Shepherd, J.G. 1997. Prediction of year-class strength by calibration regression analysis of

769 multiple recruit index series. ICES Journal of Marine Science 54: 741-752.

770

771 Shepherd, J.G. 1999. Extended survivors analysis: An improved method for the analysis of

772 catch-at-age data and abundance indices. ICES Journal of Marine Science 56: 584-591.

773

774 Sokal, R.R. and Rohlf, F.J. 1995. Biometry. Freeman, New York.

775

776 Stige, L.C., Hunsicker, M.E., Bailey, K.M., Yaragina, N.A. and Hunt G.L. 2013. Predicting fish

777 recruitment from juvenile abundance and environmental indices. Marine Ecology Progress

778 Series 480: 245-261.

779

780 Szuwalski, C.S., Vert-Pre, K.A., Punt, A.E., Branch, T.A. and Hilborn, R., 2015. Examining

781 common assumptions about recruitment: a meta-analysis of recruitment dynamics for

782 worldwide marine fisheries. Fish and Fisheries 16: 633-648.

783

784 van der Veer, H.W., 1986. Immigration, settlement, and density-dependent mortality of a larval

785 and early postlarval 0-group plaice (*Pleuronectes platessa*) population in the western Wadden

786 Sea. Marine Ecology Progress Series 29, 223–236.

787

788 Van de Wolfshaar, K.E., HilleRisLambers, R. and Gardmark, A. 2011. Effect of habitat

789 productivity and exploitation on populations with complex life cycles. Marine Ecology

790 Progress Series 438: 175-184.

791

792 Wennhage, H., 2002. Vulnerability of newly settled plaice (*Pleuronectes platessa* L.) to
793 predation: effects of habitat structure and predator functional response. Journal of
794 Experimental. Marine Biology and Ecology 269, 129–145

795

796 Zhang, T., Bailey, K.M. and Chan, K.S., 2010. Recruitment forecast models for walleye pollock
797 *Theragra chalcogramma* fine-tuned from juvenile survey data, predator abundance and
798 environmental phase shifts. Marine Ecology Progress Series 417: 237-248.

799

800 **Tables legends**

801 Table 1: The 18 species assessed by ICES in 2017-2018 whose juveniles rely on coastal
 802 habitats, and their general vertical habitat use (after Seitz et al., 2014 and updated in Brown et
 803 al., 2018a).

804

805 Table 2: The number of species and stocks assessed by ICES in 2017-2018 based on progressive
 806 sub-setting: coastal-dependent, use short-term recruitment forecasts in assessment, existence of
 807 surveys with possible estimate of pre-recruitment, and use the survey values as the predictor of
 808 recruitment in the assessment.

809

810 Table 3: The reasons for rejection, and spatial scale of the survey for the 21 stocks of species
 811 that rely on coastal habitats and for which survey-based pre-recruit abundance indices exist but
 812 are not presently used in short-term forecasts in ICES assessment.

813

814

815 Table 4: Characteristics of the 14 stocks of species relying on coastal habitats at juvenile stage,
 816 for which survey-based pre-recruit abundance indices are used in short-term forecasts in ICES
 817 stock assessments. Characteristics shown are: description of the stock, name and information
 818 on survey design (*: the selected survey indices for the 2 stocks for which 2 were available),
 819 age of pre-recruit in survey-based abundance indices, youngest age in the associated stock
 820 assessment, length of the time series, assessment model used, whether the pre-recruit survey-
 821 based indices were used in the stock assessment or only for short-term forecasts, the method to
 822 eliminate the last two years of the recruitment estimates (either “manually” or “natural, i.e.,
 823 natural elimination because the last two years were dropped when matching the two recruitment
 824 indices”), value of the correlation coefficients r_1 and r_2 (r_2 : **rerun models** (in bold) and *stocks*

825 *for which survey-indices are not incorporated in the assessment (in italic)).*

826

827

828

829

830 Figure legends

831

832 Figure 1: Number of stocks by DLS Category that used short-term forecasted recruitment in
833 their assessment, categorized by whether a pre-recruit survey exists or not, and if it exists,
834 whether it was used to predict recruitment. A total of 49 stocks were used that were species that
835 rely on coastal habitats and for which ICES assessments used short-term forecasted recruitment.
836

837 Figure 2: Scatter plot of survey-based (x axis) and assessment-based (y axis) recruitment (both
838 in the unit used in the stock assessment WG) for the 14 coastal-dependent stocks for which
839 survey-based pre-recruit abundance indices are used as short-term forecasts of recruitment in
840 ICES assessments. Stock codes are defined in Table 4.

841

842 Figure 3: Box plot of the correlation coefficients between model-based recruitment indices and
843 (left panel) the geometric mean of the model-based recruitment indices during the last five years
844 (r_3), and (right panel) the survey-based pre-recruit abundance indices (r_2). Each plot is based on
845 the 10 stocks that rely on coastal habitats at juvenile stage and for which the ICES assessments
846 are truly independent from survey-based pre-recruit abundance indices but use these survey-
847 based pre-recruit abundance indices for short-term forecasts of recruitment (thick line, median;
848 box, from the 0.25 quartile to the 0.75 quartile; whiskers, 1.5 times the distance between the
849 quartiles).

850

851

Table 1: The 18 species assessed by ICES in 2017-2018 whose juveniles rely on coastal habitats, and their general vertical habitat use (after Seitz et al., 2014 and updated in Brown et al., 2018a).

Species	Vertical position
Ammodytes	Demersal
Anguilla anguilla	Demersal
Clupea harengus	Pelagic
Dicentrarchus labrax	Demersal
Engraulis encrasicolus	Pelagic
Gadus morhua	Demersal
Limanda limanda	Benthic
Merlangius merlangus	Demersal
Mullus surmuletus	Demersal
Platichthys flesus	Benthic
Pleuronectes platessa	Benthic
Pollachius pollachius	Demersal
Pollachius virens	Demersal
Psetta maxima (historic name)	Benthic
Scomber scombrus	Pelagic
Scophthalmus rhombus	Benthic
Solea solea	Benthic
Sprattus sprattus	Pelagic

Table 2: The number of species and stocks assessed by ICES in 2017-2018 based on progressive sub-setting: coastal-dependent, use short-term recruitment forecasts in assessment, existence of surveys with possible estimate of pre-recruitment, and use the survey values as the predictor of recruitment in the assessment.

Category	Number of species	Number of stocks
ICES evaluated	61	185
& coastally-dependent juveniles	18	78
& with short-term forecast		49
& with potential existing survey based pre-recruit indices		35
& using survey-based indices in forecast		14

Table 3: The reasons for rejection, and spatial scale of the survey for the 21 stocks of species that rely on coastal habitats and for which survey-based pre-recruit abundance indices exist but are not presently used in short-term forecasts in ICES assessment.

Reason to reject	Number of stocks	Scale of the survey
Incomplete time-series	2	
Too late to be used	4	
Not investigated, nor tested	11	Stock scale, not including nurseries (2) Stock distribution partially covered, including coastal nurseries (6) Stock distribution partially covered, not including coastal nurseries (3)
Investigated and rejected	4	Stock distribution partially covered, including coastal nurseries (2) Stock distribution partially covered, not including coastal nurseries (2)

Stock Description	Stock code	area of juvenile survey	Survey name	Method of survey	Nb samples	age group of the recruitment indice	youngest age group in the stock assessment	Length of the time series	Assessment method	Incorporated in assessment and not in forecast only
Anchovy (<i>Engraulis encrasicolus</i>) in Subarea VIII (Bay of Biscay)	ane.27.8	Stock scale, including nurseries	Juvena	Accoustic	80	0	1	15	Specific SAM like	Yes
Cod (<i>Gadus morhua</i>) in Division Va (Iceland grounds)	cod.27.5a	Stock scale, not including nurseries	SMH* and (SMB)	Trawl	800	1	3	21	specific XSA like	No
Cod (<i>Gadus morhua</i>) in NAFO Subarea 1, inshore (Inshore west Greenland cod)	cod.21.1	Stock distribution partially covered, including nurseries	West Greenland inshore gill-net survey	Net	100	1	1	28	SAM	Yes
Cod (<i>Gadus morhua</i>) in Subarea IV and Divisions VIIId and IIIa West (North Sea, Eastern English Channel, Skagerrak)	cod.27.47d20	Stock scale, including nurseries	IBTS–Q1 + IBTS–Q3 combined	Trawl	200	1	1	35	SAM	Yes
Cod (<i>Gadus morhua</i>) in Subdivisions 22–24 (Western Baltic Sea)	cod.27.22-24	Stock scale, including nurseries	BITSQ4	Trawl	100	0	1	17	SAM	Yes
Herring in Subarea IV and Divisions IIIa and VIIId (North Sea autumn spawners)	her.27.3a47d	Stock scale, including nurseries	IBTS (mik)	Trawl	567	0	0	27	FLSAM	Yes
Herring in Subdivisions 25 - 29 (excluding Gulf of Riga) and 32	her.27.25-2932	Stock scale, not including nurseries	BIAS	Accoustic	49	0	1	24	XSA	No
Mackerel in the Northeast Atlantic (combined Southern, Western and North Sea spawning components)	mac.27.nea	Stock scale, including nurseries	IBTS	Trawl	1820	0	0	18	SAM	Yes
Plaice in Division VIIa (Irish Sea)	ple.27.7a	Stock scale, not including nurseries	BTS combined	Trawl	58	1	1	24	SAM	Yes
Plaice Subarea IV (North Sea)	ple.27.420	Stock distribution partially covered, including nurseries	UKBTSQ4	Trawl	100	1	1	22	AAP	Yes

Sole in Subarea IV (North Sea)	sol.27.4	Stock distribution partially covered, including nurseries	DFS combined	Trawl	630	0	1	26	AAP	No
Sprat in Subdivisions 22 - 32 (Baltic Sea)	spr.27.22-32	Stock scale, including nurseries	BIAS	Accoustic		0	1	23	XSA	No
Whiting in ICES Division VIIb, c, e-k	whg.27.7b-ce-k	Stock distribution partially covered, including nurseries	IGFS+EVHOE Combined indice	Trawl	180	0	0	14	XSA	Yes
Whiting Subarea IV (North Sea) and Division VIId (Eastern Channel)	whg.27.47d	Stock scale, including nurseries	IBTSQ3* and (IBTSQ1)	Trawl	310	1	1*	26	XSA	Yes

2 last years removed	Value of correlation coefficient (r1)	Value of correlation coefficient without survey-based index in stock assessment (r2)
	0.7	
	0.8	0.8
Manually	0.62	
Manually	0.91	
Manually	0.89	0.7
Manually	0.94	0.84
	0.92	0.92
Natural	0.64	0.58
Natural	0.67	
Manually	0.77	0.77

	0.83	<i>0.83</i>
	0.85	<i>0.85</i>
Natural	0.79	0.68
Natural	0.67	0.67




Figure 1: Number of stocks by DLS Category that used short-term forecasted recruitment in their assessment, categorized by whether a pre-recruit survey exists or not, and if it exists, whether it was used to predict recruitment. A total of 49 stocks were used that were species that rely on coastal habitats and for which ICES assessments used short-term forecasted recruitment.




Figure 2: Scatter plot of survey-based (x axis) and assessment-based (y axis) recruitment (both in the unit used in the stock assessment WG) for the 14 coastal-dependent stocks for which survey-based pre-recruit abundance indices are used as short-term forecasts of recruitment in ICES assessments. Stock codes are defined in Table 4.




Figure 3: Box plot of the correlation coefficients between model-based recruitment indices and (left panel) the geometric mean of the model-based recruitment indices during the last five years (r_3), and (right panel) the survey-based pre-recruit abundance indices (r_2). Each plot is based on the 10 stocks that rely on coastal habitats at juvenile stage and for which the ICES assessments are truly independent from survey-based pre-recruit abundance indices but use these survey-based pre-recruit abundance indices for short-term forecasts of recruitment (thick line, median; box, from the 0.25 quartile to the 0.75 quartile; whiskers, 1.5 times the distance between the quartiles).