

# Modelling the distribution of rare and data-poor diadromous fish at sea for protected area management

**Submitted to Progress in Oceanography - This version has been accepted 07/11/22)**

Sophie A. M. Elliott<sup>1, 4, 6\*</sup>, Anthony Acou<sup>1,2</sup>, Laurent Beaulaton<sup>1,5</sup>, Jérôme Guitton<sup>4</sup>, Elodie Réveillac<sup>3</sup>, Etienne Rivot<sup>1,4</sup>

<sup>1</sup> Management of Diadromous Fish in their Environment OFB-INRAE-Institut Agro-UPPA, 35042 Rennes, France

<sup>2</sup> UMS OFB-CNRS-MNHN PatriNat, Station marine du Museum National d'Histoire Naturelle, 35800 Dinard, France

<sup>3</sup> Littoral, Environnement et Sociétés (LIENSs), UMR 7266, La Rochelle Université-CNRS, 17000 La Rochelle, France

<sup>4</sup> DECOD (Ecosystem Dynamics and Sustainability), L'Institut Agro, Ifremer, INRAE, Rennes, France

<sup>5</sup> Service Conservation et Gestion Durable des Espèces Exploitées, OFB, DRAS, 35042 Rennes, France

<sup>6</sup> Game & Wildlife Conservation Trust, Salmon & Trout Research Centre, East Stoke, Wareham BH20 6BB, UK

\*Corresponding author: E-mail address: SElliott@gwct.org.uk (S.A.M. Elliott)

## ABSTRACT

Anthropogenic pressures have resulted in declines in diadromous fish. Many diadromous fish which were commercially important are now threatened and protected. Little is known about their marine life history phases, and no observation-based Species Distribution Model exists for this group of species at sea. Yet, fisheries dependent and independent data could provide new insights into the distribution of diadromous fish at sea.

We collated a database of 168 904 hauls from fisheries observer bycatch data and scientific fisheries surveys, from eastern Atlantic and Mediterranean waters. The distribution of eleven rare and data-poor diadromous fish (shads, lampreys, salmonids, the European eel, the thinlip mullet, smelt and the European flounder) were modelled. A Bayesian site occupancy model, that incorporates imperfect detection to account for repeat detections and non-detections, the non-random nature of fishing gear type and spatial autocorrelation was used. From the model outputs, we explored bycatch risk and the role of MPAs, required under the Marine Strategy Framework Directive and Habitat Directive and assessed.

Diadromous fish were observed within relatively shallow coastal areas. Species specific gear bycatch trends were observed. Core distribution areas corresponded to their known water basin presence, indicating connectivity with their freshwater habitats. Numerous Habitat Directive Marine Protected Areas were found to be of relevance.

Given the coastal distribution of these species, they are exposed to higher anthropogenic pressures from both terrestrial and marine environments. Risk of bycatch at sea for most species appears to be low. Nonetheless, for threatened individuals, even a small amount of bycatch may impact their populations, especially since misreporting is likely to be high. Differences in catchability between gears highlight potential benefits of limiting access of certain gears within protected areas to reduce bycatch.

**Key words:** Species Distribution Model, diadromous fish, rare species, imperfect detection, Marine Protected Areas, bycatch.

# 1. Introduction

Diadromous fish undertake long distance migrations between freshwater and marine ecosystems during their varying life history stages (Lassalle et al., 2008; Legrand et al., 2020; Limburg and Waldman, 2009; McDowall, 2009). They are particularly vulnerable since they are subject to both terrestrial and freshwater pressures (e.g., terrestrial run-off, pollution, habitat destruction, barriers to migration, climate change, fishing, etc.) (Costa et al., 2021; Limburg and Waldman, 2009; Merg et al., 2020; Verhelst et al., 2021). Furthermore, unlike many marine fish that are geographically widespread, numerous anadromous fish (e.g., shad and salmonids) form river-specific populations that are more susceptible to extinction (Limburg and Waldman, 2009; McDowall, 2009). As a result, most diadromous species native to the northern Atlantic Ocean have declined in abundance by at least 90% since the end of the late 19<sup>th</sup> century (Drouineau et al., 2018; Limburg and Waldman, 2009; Waldman and Quinn, 2022).

To try and halt the loss of diadromous fish populations, various International Agreements (e.g., Bern Convention, Convention of Migratory Species) and legislations (e.g., EU Habitat Directive, EU Marine Strategy Framework Directive (MSFD)) have been enacted (Table 1). Measures implemented focus principally on freshwater or estuarine habitats, such as restoration of river continuity as required by the EU Water Framework Directive (WFD, 2000/60/EC) and freshwater Spatial Areas of Conservation under the Habitat Directive (HD, 92/43/EEC). However, most diadromous species remain threatened at a national level ([www.nationalredlist.org](http://www.nationalredlist.org); Table 1), and the improvements have yet to be attained throughout their range (Verhelst et al., 2021; Waldman and Quinn, 2022; Wilson and Veneranta, 2019). The marine life history stages and at-sea distribution of diadromous fish, largely remain a black box (Elliott et al., 2021), causing difficulty in evaluating their conservation status required under the various legislations that protect these species (Table 1; Wilson and Veneranta, 2019). Furthermore, few Marine Protected Areas (MPAs) have been designated to protect them (e.g., <https://mpa.ospar.org>; <https://natura2000.eea.europa.eu>), and at-sea bycatch is not well understood, even if it is suspected that it could contribute to significant mortality (Kappel, 2005; Stratoudakis et al., 2016; Verhelst et al., 2021; Wilson and Veneranta, 2019).

Under the EU data collection framework, Member States are required to collect fisheries bycatch data through an onboard observer program (Cornou et al., 2015). Fisheries data is a rich set of information which provides year-round catch information and can be a valuable source of information for data-deficient species (e.g., Baum et al., 2003; Bisch et al., 2022; Elliott et al., 2020b). Bycatch data for protected species can be difficult to access, therefore fisheries observer data could help understand the distribution and bycatch mortality of data-poor species and meet EU Habitat Directive and MSFD (92/43/EEC; 2008/56/EC) requirements. Use of fisheries data does, however, require biases from the different gear types, the targeted nature of fishing, and un-balanced sampling to be taken into consideration (e.g. Alglave et al., 2022; Bourdaud et al., 2017).

Species Distribution Models (SDMs) provide a means to ‘fill in the gaps’ and provide complete coverage maps on which to base conservation and management decisions (Guisan and Thuiller, 2005; Guisan and Zimmermann, 2000; Leathwick et al.,

2005). Modelling the distribution of diadromous fish could help improve protection measures for threatened and data-poor (insufficient biological information to determine the current exploitation status (Berkson and Thorson, 2015; Prince and Hordyk, 2019)) diadromous fish within existing MPAs. SDMs could also be used to meet Descriptor 1 (Biodiversity - habitat extent) requirements under the MSFD (2008/56/EC). However, building diadromous fish at sea distribution models remains challenging because most of these fish are either IUCN red listed at a national level ([www.nationalredlist.org](http://www.nationalredlist.org)) or data-poor (Limburg and Waldman, 2009; Merg et al., 2020). Modelling the distribution of data-poor and rare species, can lead to underestimation of their true distribution due to imperfect detection (uncertainty in the presence of a species within a site where the data was recorded) (Guillera-Arroita, 2017; MacKenzie et al., 2002).

To model rare and data-poor species' distribution, specific statistical methods are required to combine several sources of data that may not be initially designed to study the species distribution (Engler et al., 2004; Lomba et al., 2010; Simmonds et al., 2020). Using such data requires correction for detection bias arising from observer error, species rarity and variability in environmental conditions (Guillera-Arroita, 2017; Kellner and Swihart, 2014; MacKenzie et al., 2002) to help determine sites where the probability of presence of species is higher (Belmont et al., 2022). The low detectability of rare species can also result in a high proportion of false absences (Guillera-Arroita, 2017). Spatial and temporal variation in observation effort is another important source of heterogeneity in the observations (Guillera-Arroita, 2017; Kellner and Swihart, 2014; Meyer et al., 2011). Integrated hierarchical statistical model for species distributions have a great potential to combine multiple sources of data to infer a single latent field of abundance or presence/absence, and to enhance rare species distribution models for conservation and management purposes (Engler et al., 2004; Lomba et al., 2010; Simmonds et al., 2020). Another common problem which can arise when modelling species distribution is that data can be autocorrelated (more similar when closer together). Residual spatial autocorrelation can arise from population demographic processes or the influence of unobserved variables. Ignoring spatial autocorrelation can lead to inaccurate parameter estimations (Latimer et al., 2006). Developing models that explicitly account for spatial autocorrelation in the latent field of presence/absence is, therefore, necessary for accurate inferences on the distribution.

Given the lack of knowledge on the distribution of most diadromous fish during their marine life history stages, and the need to ensure they are sufficiently protected, our objective was to model the current distribution (2003-2019) of diadromous fish at sea. In addition, we wanted to quantify the risk of bycatch from different gear types and assess the value of Habitat Directive MPAs for their conservation. Since at sea targeted surveys do not exist for these species, we collated a database of 168 904 hauls from fisheries dependent (French fisheries bycatch observer data) and independent data (scientific survey data undertaken to assess fish populations) in eastern Atlantic and French Mediterranean waters. A Bayesian site-occupancy model, taking into account spatial autocorrelation in the probability of presence, and imperfect detection through gear type, was used for these data-poor species (Dormann et al., 2007; Latimer et al., 2006; MacKenzie et al., 2002; Moriarty et al., 2020). From the distribution models we wanted to

161 answer the following questions: 1) Is the distribution of diadromous fish at sea  
162 connected to existing knowledge on their freshwater habitats? 2) Do different gears  
163 have different catchability, and does this provide information on the risk of  
164 bycatch of diadromous fishes? 3) Could existing Habitat Directive MPAs protect  
165 diadromous fish if specific measures are implemented?

166 **Table 1.** Diadromous fish observed within north-eastern Atlantic waters and their IUCN conservation status. A = Anadromous, C =  
167 Catadromous, - = not listed, CR = Critical, LC = Least Concern, VU = Vulnerable. Demersal species are found near the seabed,  
168 pelagic species occur in the open sea, demerso-pelagic species are species that migrate between these two water column zones,  
169 host dependent refers to lampreys that are parasitic species during their marine phase. ✓ = protected under the specific  
170 convention or Directive, letters specify the specific appendix the species is protected under.

Latin name	Common name	Type	Water column zone <sup>1</sup>	EU IUCN <sup>3</sup>	International conventions and legislation							
					CITES	WFD	HD	MSFD	Bern	Bonn	OSPAR	Barcelona
<i>Acipenser sturio</i>	Atlantic sturgeon	A	Demersal	CR	AI	✓	II, IV	✓	III	I, II	✓	✓
<i>Alosa alosa</i>	Allis shad	A	Pelagic	LC	-	✓	II, V	✓	III	-	✓	-
<i>Alosa fallax</i>	Twait shad	A	Pelagic	LC	-	✓	II, V	✓	III	-	-	-
<i>Alosa agone</i>	Mediterranean twaite shad	A	Pelagic	LC	-		II, V	✓	III	-	-	✓
<i>Anguilla anguilla</i>	European eel	C	Demerso-pelagic	CR	All	✓	-	✓	-	-	✓	✓
<i>Lampetra fluviatilis</i>	River lamprey	A	Host dependent	LC	NA	✓	II, V	✓	III	-	-	✓
<i>Petromyzon marinus</i>	Sea lamprey	A	Host dependent	LC	-	✓	II	✓	III	-	✓	✓
<i>Chelon ramada</i> *	Thinlip mullet	C	Demersal <sup>2</sup>	LC	-	✓	-	-	-	-	-	-
<i>Osmerus eperlanus</i>	Smelt	A	Pelagic	LC	-		-	-	-	-	-	-
<i>Platichthys flesus</i>	European flounder	C	Demersal	LC	-		-	-	-	-	-	-
<i>Salmo salar</i>	Atlantic salmon	A	Pelagic	VU <sup>4</sup>	-	✓	II, V	✓	III	-	✓	-
<i>Salmo trutta</i>	Sea trout	A	Pelagic	LC	-	✓	-		-	-	-	-

171   <sup>1</sup> Elliott and Dewailly, 1995;<sup>2</sup> Almeida, 1996; <sup>3</sup> Freyhof and Brooks, 2011; <sup>4</sup> Nieto et al., 2015 \*previously known as *Lisa ramada*.

## 2. Method

### 2.1. Combining fisheries independent and dependent surveys

Fisheries dependent and independent data from 1965 to 2019 were collated within eastern Atlantic waters (Greater North Sea, Celtic Sea, Bay of Biscay and the Iberian coast) and French Mediterranean waters). Scientific bottom trawl surveys were extracted from the International Council for the Exploration of the Sea (ICES) Database of Trawl Surveys (DATRAS) portal (<https://www.ices.dk/data/data-portals/Pages/DATRAS.aspx>; Refer to Elliott et al., 2021 Table S1; Data in brief Table S1 - S2) and French Metropolitan scientific surveys not submitted to ICES DATRAS were obtained from the Institut Français de Recherche pour l'Exploitation de la Mer (IFREMER) (<https://campagnes.flotteoceanographique.fr/campaign>). Fisheries-dependent data came from French fisheries observer data (ObsMer data; Cornou et al., 2015) that began in 2003. According to the sampling plan, fisheries observers sample fishing vessels and fishing operations when on board (Fauconnet et al., 2015). ObsMer data is held by IFREMER and available on request from the French Ministry of Fisheries and Aquaculture (Direction des Pêches Maritimes et de l'Aquaculture; Refer to Elliott et al., 2021 Table S1; Data in brief Table S1).

From the database, diadromous fish presence and absence, length, gear type, spatial location and year and month of capture were extracted (Fig. 1; Fig. S1). Due to missing data and insufficient information on the length of hauls, size of vessels, mesh size, etc., effort was not possible to calculate. Details of gear types and surveys that were used for the analysis can be found within the supplementary materials of Elliott et al., (2021).

Prior to model analysis significant data cleaning was undertaken to standardise the surveys and extract relevant data to perform statistical analysis (e.g., converting gear types to gear categories, converting alike variables to the same units, etc.; see details within the Data in brief; Appendix A). Data from 2003 were used for spatial analysis, since a large proportion (74%) of the data came from the ObsMer dataset which started in 2003. In addition, prior to this data few presences were observed.

Since the proportion of zeros to presences was very uneven, which is not ideal for predictive modelling (Fielding and Bell, 1997), all surveys, gear types, target species (the intended catch for a particular fishery - from the ObsMer dataset) and ICES statistical divisions (Fig. 1) without diadromous fish presence, were deleted before modelling (Fig. S2 contains a spatial map of all hauls per gear type). Depths outside the ranges the species were observed were also removed. To avoid targeted fishing bias, all species that were targeted were removed. The latter included 2 European eels, *Anguilla anguilla*, 33 thinlip mullets, *Chelon ramada* and 41 flat fish which may have included European flounders, *Platichthys flesus* presences. All species were modelled other than the European sturgeon, *Acipenser sturio* because of too few presences (11 presences from 2003 - 2019). Note, for *L. fluviatilis*, one individual was caught by Set Longline within the Bay of Biscay. Due to its isolated nature of this capture, this caused problems when modelling (over detection of the line caught individual). Line gear type was therefore removed from the hierarchical modelling process.



## 2.2. Environmental predictor variables

Six environmental variables were considered as potential predictors of the presence of diadromous fish (seabed depth; distance from coast; sediment type; salinity; net primary production; and sea surface temperature; Table 2; Fig. S3). Depth and distance from the coast are thought to indicate key diadromous fish migratory periods (Taverny et al., 2012; Taverny and Elie, 2001; Trancart et al., 2014). Salinity and temperature are known to have direct physiological effects on diadromous fish, and their changes can indicate migration timing (Arevalo et al., 2020; Trancart et al., 2014). Net primary production is the gross primary production by autotrophs over the rate at which they respire, which is a measure of marine ecosystem functioning (ERSEM, 2020). Sediment type can be considered a proxy for food availability and shelter (Elliott et al., 2016; Trancart et al., 2014). As of result of too few presences and less information on length than the presence of individuals, distribution changes in length at stage, seasonal life history stage migration, and changes in habitat occupancy over the years were not considered.


To select the model of best fit, variables that were not collinear (Pearson's correlation coefficient  $< 0.5$  or  $> -0.5$  and a Variance Inflation Factor  $< 2$ ) (Dormann et al., 2013) were kept. Depth and distance from coast, and sea surface temperature and net primary production were collinear. Since most species occupied a wide latitudinal range and seasonal or yearly temperature effects were not modelled, net primary production was used instead of sea surface temperature. To identify the model of best fit a backwards stepwise model selection process was undertaken, ensuring distance from coast and depth were not in the same model. The model with the lowest deviance information criterion, which includes a penalty factor for the number of parameters, was selected (Spiegelhalter et al., 2002).

**Table 2.** Environmental predictors used to analyse diadromous fish distribution. For all species except *Alosa agone*, variables were extracted at a 0.1 x 0.1 decimal degree resolution. *A. agone* environmental variables were extracted at a 0.04 decimal degree resolution.

Variable	Unit	Description / Source
Depth	Meters (m)	Bathymetry data downloaded from National Aeronautics and Space Administration Earth Observations project ( <a href="http://neo.sci.gsfc.nasa.gov">http://neo.sci.gsfc.nasa.gov</a> ) with a 0.1 x 0.1 grid cell resolution by using the General Bathymetric Chart of the Oceans grid.
Distance to coast	Kilometres (km)	Calculated from a continent shape file
Sediment	Categorical	Broad-scale seabed habitat types (EUSeaMap) downloaded from EMODnet. Four classes were used according to grain size: mud ( $\leq 2$ mm), sand ( $> 2$ and $\leq 4$ mm), coarse grain ( $> 4$ mm and $\leq 64$ mm), and rock ( $> 64$ mm).
Salinity	Practical Salinity Units (PSU%)	Modelled sea surface salinity at 0.1 x 0.1 decimal degree resolution downloaded from Copernicus (ERSEM, 2020).

Net primary production	Milligrams of carbon per meters squared (mg C/m <sup>2</sup> )	Modelled at 0.04 x 0.04 decimal degree resolution from Copernicus using Teruzzi et al., (2021). Modelled net primary production at 0.1 x 0.1 decimal degree resolution downloaded from Copernicus (ERSEM, 2020). Modelled at 0.04 x 0.04 decimal degree resolution from Copernicus using Teruzzi et al., (2021).
Sea surface temperature	degrees centigrade (°C)	Modelled Sea surface temperature at 0.1 x 0.1 decimal degree resolution downloaded from Copernicus (ERSEM, 2020). Modelled at 0.04 x 0.04 decimal degree resolution from Copernicus using Teruzzi et al., (2021).

253



**Fig. 1.** Diadromous fish presence (blue dots) and absence (light grey dots) data used to model their distribution (2003-2019). Dark grey lines indicate International Council for the Exploration of the Sea statistical divisions.

### 2.3. Statistical analysis

A Bayesian hierarchical site occupancy Gaussian intrinsic Conditional Autoregressive (iCAR) model was built independently for each species. Hierarchical models have the advantages of separating the ecological process (here, the habitat suitability model, that is the spatial distribution of the probability of presence/absence, including spatial autocorrelation) from the observation process (here, the imperfect detection and its variability among gears or other factors) (Dormann et al., 2007; Isaac et al., 2020; Latimer et al., 2006; MacKenzie et al., 2002). The fundamental concept of the model is that the true presence/absence spatial field (including spatial autocorrelation) is modelled on a grid, each grid cell being associated with multiple observation events with imperfect detection, with detectability that may vary depending upon the fishing gear.

The site occupancy model integrates two processes:

#### i) Latent ecological process (habitat suitability)

The latent field of probability of presence is modelled on a grid as a function of environmental predictors and by explicitly accounting for spatial autocorrelation:

$$\text{logit}(\theta_j) = X_j\beta + P_j \quad (1)$$

with  $\theta_j$  the latent probability of presence (the habitat suitability) within grid cell  $j$ , modelled in the logit scale as a function of environmental predictors  $X_j$ , with  $\beta$  a vector of fixed effect describing how much the environmental predictor contribute to the suitability process. The spatial autocorrelation  $P_j$  is modelled at the scale of the grid cell as an iCAR random effect (2):

$$\text{iCAR: } P_j \sim \text{Normal}(\mu_j, \frac{V_p}{n_j}) \quad (2)$$

with  $\mu_j$  the mean of the random effects calculated over the  $n_j$  cells considered in the neighbour of cell  $j$  ( $n_j$  is the number of the neighbours which is 8 in most cases, and  $<8$  when the cell is on the boundary of the spatial domain),  $V_p$  the variance of the spatial random effect. A site is defined here as the area/volume covered during a given sampling/fishing operation. Any site  $i$  within the cell  $j$  as the same latent probability of presence  $\theta_j$ .

#### ii) Observation process (detection):

Conditionally upon the latent probability of presence/absence in each grid cell  $j$  as describe above, the multiple detection events  $y_{j,i}$  (data 0/1) associated to the same grid cell  $j$  are modelled as mutually independent two steps Bernoulli process with detectability  $\delta_{j,i}$ :

$$y_{j,i} \sim \text{Bernoulli}(z_{j,i} \delta_{j,i}) \quad (2)$$

with

$$z_{j,i} \sim \text{Bernoulli}(\theta_j) \quad (3)$$

and

$$\text{logit}(\delta_{j,i}) = W_{j,i}\gamma$$

where  $\delta_{j,i}$  denotes the probability of detecting the species at site  $i$  within cell  $j$ , modelled as a fixed effect of the gear type associated with observation  $i$ .  $W_{j,i}$  is the raw vector of the design matrix, and  $\gamma$  is the vector of the gear effects, so as  $W_{j,i}\gamma$  is the gear affect associated with observation at site  $i$ .

All SDMs were undertaken using the ‘hSDM’ package (Vieilledent et al., 2014) using the ‘mod.hSDM.siteocc.iCAR’ function in R version 3.6.3 (R Code Team 2018). The prior distribution for the variance of the spatial random effect followed weakly informative uniform distribution. The effect of the detectability from gear and from the suitability process was modelled as fixed effects drawn from an weakly informative Gaussian prior centred at zero with a fixed standard deviation of 2 (Gelman et al., 2008; Northrup and Gerber, 2018). Each model was run with 50 000 Gibbs iterations, five Markov chain Monte Carlo simulations and a burn-in phase of 50 000 iterations.

#### 2.4. Grid size and spatial distribution

A critical issue in our modelling framework is the choice of the grid size. Since the latent field of presence/absence is modelled at the scale of the grid cell (eq. 1), the larger the grid resolution the higher the probability to get at least one presence within a grid cell. A finer grid cell will limit the quantity of available observations falling within each cell, which therefore increases difficulty in model fitting. We selected the finest grid as possible that allow the model to converge. For almost all species modelled within Eastern Atlantic waters, a grid of 0.1x0.1 decimal degrees was used. Because of the scarcity of data for Lampreys, the model did not converge with such a resolution and a coarser resolution of 0.2x0.2 decimal degrees was therefore used for these species. For *A. agone*, that only occurs within the Mediterranean, a grid resolution of 0.04 x 0.04 decimal degrees was used because of the smaller area encompassed. This spatial resolution (0.04 x 0.04 decimal degrees) was not used for other species since it was not available at the scale required within Eastern Atlantic waters. All environmental variables were then aggregated and modelled to their relevant spatial resolution. There were insufficient presences within the Mediterranean for *A. anguilla*, *P. marinus*, *C. ramada*, *P. flesus* predictions (issues with model convergence). Predictions within the Mediterranean for these were therefore excluded (Fig. 2).

#### 2.5. Model evaluation

The probability of presence was calculated from the mean of the posterior distribution of the latent field  $\theta_j$ . A threshold of 0.4 was used to bin estimated probability of presence into binary presence/absence values. A threshold of 0.4 was used as a compromise given the rarity of the species and accuracy following examining various thresholds and plotting positive predicted values. The model performance was then evaluated using sensitivity (proportion of correctly classified presences), specificity (proportion of correctly classified absences) and percent correct classification (proportion of presences and absences correctly using the R package ‘PresenceAbsence’ (Freeman and Moisen, 2008a; Fielding and Bell, 1997; Table 3). Max sensitivity specificity which is often used for rare species distribution models (Freeman and Moisen, 2008b; Jiménez-Valverde and Lobo, 2007), was not used since false presences was very high when applying this metric. Additionally, our SDM objective was to try and minimise false presences and absences within the distribution models. The threshold independent Area under the receiver operating curve was also used to evaluate the model performance. These four metrics were used to better understand the accuracy of the model since, some metrics can be biased with low prevalence data and where imperfect detection can arise (Allouche et al., 2006; Fourcade et al., 2018; Guillera-Arroita, 2017; Leroy et al., 2018).



So that areas with probabilities of presence above the threshold could be easily identified for management purposes, all predictions below 0.4 were mapped with a uniform light blue colour (Fig. 2). Uncertainty maps for each model were also produced to gauge a better idea of areas of prediction with higher uncertainty (Loiselle et al., 2003; Spiegelhalter et al., 2002). The uncertainty maps were calculated by the mean coefficient of variation from the posterior distribution of spatial parameters in the model of best fit for each species (Lambert et al., 2020; Latimer et al., 2006). Probabilities of prediction and uncertainty above 0.4 were classified into 0.2 units to highlight areas of higher and lower probability of presence more clearly (Fig. 2; Fig. S4). As a result of the low diadromous fish presence, splitting the data into learning/validation samples led to insufficient presence in the learning data set. Cross validation was therefore not undertaken.

## 2.6. Diadromous fish river basin presence

The predicted distributions were visually compared to the distribution of diadromous fish within their freshwater habitats using the EuroDiad database (version 4; <https://data.inrae.fr/dataset.xhtml?persistentId=doi:10.15454/IVVAIC>; Barber-O'Malley et al., 2022). From EuroDiad database, migrating diadromous fish population presence within the study areas our model encompassed, were extracted using 'recent' literature citing from 1951 to 2020 (see Barber-O'Malley et al., (2022) and Béguer et al., (2007) for more details on the database) and mapped with the hierarchical SDM results (Fig. 2). The EuroDiad4 database was used as opposed to IUCN red list map distribution since it has been peer reviewed and it is more up to date.

## 2.7. Habitat Directive MPAs of relevance to diadromous fish

'Core areas' with a probability of presence >0.4 were selected and intersected with European Union Natura 2000 Habitat Directive Sites of Community Importance (<https://www.eea.europa.eu/data-and-maps/data/natura-12>). A threshold of > 0.4 probability of presence was selected as a result of the low probability of presence of diadromous fish (particularly Habitat Directive protected species), and to ensure consistency with the model evaluation threshold. Since only one Site of Community Importance was submitted to the European Union from the United Kingdom, Special Areas of Conservation were taken into consideration (<https://jncc.gov.uk/our-work/special-areas-of-conservation-overview/>). For each species, the number of MPAs (Sites of Community Importance and Special Areas of Conservation) and the proportion of MPAs of relevance to diadromous fish (core areas within an MPA) were recorded. The extent of the core area within the MPAs was not calculated as this would be dependent of the model grid used (Table 3). Furthermore, numerous MPAs were smaller than the grid.

# 3. Results

## 3.1. Diadromous fish hierarchical SDM

For all species the Area under the receiver operating curve and Percent Correct Classification performed very well ( $\geq 75\%$ ; Table 3). Correct presence classification from the confusion matrices was consistently lower ( $< 15\%$ ) than that of correct absence classification ( $\geq 64\%$ ), and false absences were lower ( $\leq 3\%$ ) than false presences ( $\leq 22\%$ ) because of presence absence imbalance (Table 3; Table S1). The later led to on average better specificity than sensitivity scores (Table 3).

*P. flesus* was the most prevalent species modelled (5394 presences), whereas salmonids the least (*S. salar* = 68 presences and *S. trutta* 63 presences) (Table 3). SDMs that had lower presences (salmonids and lampreys) did not predict as well (lower sensitivity and higher uncertainty maps) as species with higher presence (shads, *C. ramada*, *P. flesus* and *O. eperlanus*). In addition, species that had a more disperse distribution (e.g., *A. anguilla* and *S. salar*) did not predict as well (sensitivity  $\leq 0.66$  for both species) as those with a more aggregated distribution (*O. eperlanus* sensitivity = 0.81) (Table 3; Fig. 1 and 2). Uncertainty maps showed that in general, areas of high and low predicted probability of presence had lower uncertainty (Fig. S4).


All species were observed to have a relatively shallow (<300m depth) coastal distribution (~<300 km from the coast; Fig. 2 and 3). The main predictor variables for diadromous fish habitats were depth or distance from coast and salinity (Fig. 3). Since sufficient *C. ramada* were observed within eastern Atlantic and Mediterranean waters to model, two salinity peaks are observed corresponding to the different salinity ranges of the two regions (Fig. 3).


424 **Table 3.** Hierarchical species distribution model outputs. All models contained gear as part of the observation process. AUC =  
 425 Area Under the receiver operating Curve score, PCC = Percent Correct Classification, NetPP = Net Primary Production. Max depth  
 426 is the maximum depth at which the dataset was cut. Note most environmental variables (all except distance and depth) were  
 427 not available within very coastal areas (e.g., within estuaries and inlets). Numerous presences and absences were therefore lost  
 428 when extracting the environmental variables for modelling.

Species	Model selected	Hauls	Presence	Absence (%)	Max depth (m)	Sensitivity	Specificity	AUC	PCC	Cell size (decimal degrees)
<i>A. alosa</i>	Depth + Salinity + NetPP + Sediment	86 292	802	99	300	0.79	0.92	0.95	0.95	0.1x0.1
<i>A. fallax</i>	Distance + Salinity + NetPP + Sediment	91 135	1385	98	300	0.88	0.81	0.90	0.82	0.1x0.1
<i>A. agone</i>	Depth + Salinity + Sediment	5086	176	97	200	0.81	0.74	0.86	0.75	0.04x0.04
<i>A. anguilla</i>	Depth + Salinity + NetPP	54 124	176	99	250	0.53	0.99	0.98	0.98	0.1x0.1
<i>L. fluviatilis</i>	Distance + Salinity	16 536	68	99	150	0.71	0.95	0.96	0.94	0.2x0.2
<i>P. marinus</i>	Distance + Salinity	28 582	74	99	300	0.60	0.98	0.94	0.97	0.2x0.2
<i>C. ramada</i>	Distance + Salinity + Netpp + Sediment	53 083	925	98	300	0.84	0.93	0.96	0.92	0.1x0.1
<i>P. flesus</i>	Distance + Salinity + NetPP + Sediment	92 400	5394	94	300	0.90	0.93	0.97	0.93	0.1x0.1
<i>O. eperlanus</i>	Depth + Salinity + NetPP	48 877	1035	98	450	0.81	0.99	0.99	0.99	0.1x0.1
<i>S. salar</i>	Depth + Salinity + NetPP	36 008	68	99	150	0.66	0.98	0.95	0.98	0.1x0.1
<i>S. trutta</i>	Depth + Salinity + NetPP	31 754	63	99	150	0.71	0.94	0.93	0.93	0.1x0.1


429








**Fig. 2.** Diadromous fish hierarchical species distribution model probability of presence. Darker blue shades = SDM probability of presence > 0.4, light blue shade = SDM probability of presence ≤ 0.4. Habitat Directive Marine protected Areas (MPAs) outlined in black. MPAs of relevance to core (probability of presence > 0.4) diadromous fish distribution, highlighted in dark red. Dark red cells are larger than the gridded cells to ensure smaller core areas within MPAs are visible. Darker grey terrestrial areas represent river basins which the diadromous fish migratory populations have been observed (EuroDiad V.4). Note, no data were available within the EuroDiad database from Dutch river basins.





**Fig. 3.** Diadromous fish hierarchical species distribution model environmental predictor effects. NetPP = Net Primary Production, C = Course grain, M = Mud, S = Sand, R = Rock. The solid line represents the smooth function estimate; the shaded region represents the approximate 95% credibility interval. Boxplot blue stars represent means and small black dots the predictions.

### 3.2. *Diadromous fish river basin connectivity*


2717 references contained within the EuroDiad database were used to map the 11 diadromous fish presence within the river basin. Core distributions of diadromous fish at sea largely matched with the freshwater areas of presence (Fig. 2). The spatial extent of diadromous fish at sea was often less than that observed within river basins (e.g., lampreys, *C. ramada*, *O. eperlanus* and *S. trutta*; Fig. 2). It should be noted that there are gaps in the EuroDiad database (refer to Barber-O'Malley et al., (2022)). For example, no diadromous fish presence data were available within Dutch freshwater habitats, and salmon populations are known to exist within Danish waters (de Groot et al., 2012; Maes et al., 2007; Rikardsen et al., 2021). Presences of *A. alosa* and *A. fallax* within the Mediterranean from the EuroDiad database will have been due to historic classification (Bagliniere and Elie, 2000; Keith et al., 2020).

### 3.3. *Gear type detectability*

All species were caught by a range of gear categories ranging from benthic, demersal and pelagic mobile trawls, seine nets, line gear types (Fig. 4; Table S2). Some general patterns were, however, evident. Line gear types caught the least diadromous fish (2% (weighted), 17 presences over ~2887 hauls), whereas demersal mobile trawl gear types (otter beam trawl and otter twin trawl) caught the most (67%, 7409 presences over ~82 531 hauls; Fig. 4; Table S2). Shad had a higher detectability with static and demersal mobile gears. *A. anguilla* were detected by a wide range of gear types. Lamprey were mainly detected by mobile trawl gears. *P. flesus* and *O. eperlanus* had a higher detectability with demersal and benthic mobile gear, and salmonid had a higher detectability with static nets and pelagic gears. Rarer species and species that were caught by a wider range of gear types had lower detectability than more abundance species caught by fewer gear types (Fig. 4). On average static nets (e.g., trammel net and set gillnets) and pelagic mobile gears (e.g., Otter midwater trawls) caught larger diadromous fish than mobile demersal and benthic gear types (Fig. S1). Although the modelled species were not target species of the fisheries within the observer database, some species were landed (Table 4).



482



**Fig. 4.** Matrix of the posterior mean (marginal posterior distribution) for gear A) detectability (inverse logit of  $\gamma$ ; the probability to detect the presence of fish when they are present) and B) its standard deviation. Lighter blue shades represent higher detectability (A) and darker green shades higher detectability standard deviation (B).

**Table 4.** Landed and discarded diadromous fish from the ObsMer data. Excluding targeted catches. Proportion landed = total landed / landed + discards.

Species	Landed	Discard	Hauls	Proportion landed
<i>A. alosa</i>	182	515	70 378	0.26
<i>A. fallax</i>	152	608	64 300	0.20
<i>A. agone</i>	28	69	7373	0.29
<i>A. anguilla</i>	10	8	28 194	0.56
<i>L. fluviatilis</i>	0	2	828	0.00
<i>P. marinus</i>	2	18	10 030	0.10
<i>C. ramada</i>	643	101	45 699	0.86
<i>P. flesus</i>	856	648	66 196	0.57
<i>O. eperlanus</i>	6	45	31 311	0.12
<i>S. salar</i>	29	27	26 009	0.52
<i>S. trutta</i>	43	18	31 666	0.70

### 3.4. Habitat Directive MPAs relevance

A total of 482 Habitat Directive MPAs have been designated within the area in which the species were modelled (Fig. 2). Despite the small area of core habitats relative to the model prediction area, for most Habitat Directive listed species, numerous MPAs overlap with species core habitats, indicating their potential for the protection of diadromous fish at sea (Table 5; Fig. 2).

**Table 5.** Habitat Directive Marine Protected Areas of relevance to core diadromous fish distribution (probability of presence >0.4). Diadromous fish listed under the Habitat Directive in bold.

Species	Number MPAs of relevance	Proportion of MPAs of relevance
<i>A. alosa</i>	35	0.08
<i>A. fallax</i>	60	0.13
<i>A. agone</i>	10	0.27
<i>A. anguilla</i>	19	0.04
<i>L. fluviatilis</i>	11	0.02
<i>P. marinus</i>	14	0.03
<i>C. ramada</i>	23	0.05
<i>P. flesus</i>	65	0.15
<i>O. eperlanus</i>	13	0.03
<i>S. salar</i>	17	0.04
<i>S. trutta</i>	28	0.06

#### 4. Discussion

To our knowledge this is the first comprehensive study to model the distribution of diadromous fish at sea across eastern Atlantic and French Mediterranean waters. Using a hierarchical model allowed us to integrate an extensive database that we collated, containing both fisheries dependent and independent data to model the distribution of eleven rare diadromous fish over a wide spatial area. The model also enabled us to evaluate the probability to detect the presence of fish by the different gear types, and thus provide key information to evaluate the risk of bycatch of diadromous fish by commercial fisheries at sea.

Diadromous fish are bound to coastal areas because of their migratory behaviour. They are known to be fragile species because of the physiological and osmoregulatory changes required to undertake migrations between fresh water and marine environments (McDowall, 2009). Yet, coastal ecosystems, particularly within eastern Atlantic and Mediterranean waters, are subject to some of the highest anthropogenic impacts (Halpern et al., 2015; Korpinen et al., 2021; Lotze et al., 2006). Even though more data were collected within coastal areas, and the species are known to migrate from freshwater habitats, our study confirms that the investigated species were found to mainly concentrate in coastal and relatively shallow waters. The strong dependency of diadromous fish species to coastal areas, as evidenced in this study, reinforces the importance of investigating the vulnerability of diadromous fish to coastal anthropogenic pressures (Halpern et al., 2015; Limburg and Waldman, 2009; Lotze et al., 2006; Worm et al., 2006).

##### 4.1. Hierarchical Bayesian SDM

On average, the eleven SDMs performed well. All the models had lower sensitivity and positive prediction values scores, because of low presence. Areas of both high probability and very low probability of presence had lower uncertainty, giving a



more accurate picture of confidence in the distributions (e.g., higher certainty that most species are absent within waters further from the coast). In addition to problems with very low occurrence, the areas of higher uncertainty for certain species (e.g., *S. salar* and *S. trutta*, *A. alosa* and *A. fallax*), may be partially because of misidentification between these groups of diadromous fish. The core areas of probability of presence, in combination with the confidence maps provide a better understanding of confidence of these predictions. To improve problems of false presence predictions which might occur with species that are difficult to identify, Bayesian models incorporating false positive errors could be undertaken (e.g. Diana et al., 2021; Guillera-Arroita, 2017; Royle and Link, 2006).

Separating out the spatial random field of presence absence from the observations enabled us to explicitly consider the imperfect detection of the observation process. Since our database contained non-diadromous fish specific survey and catch data which sampled the same grid cell numerous times, the use of the site occupancy model enabled imperfect detection parameters to be estimated. Disregarding false absences would have led to biased inferences and over-stated parameter precision (Guillera-Arroita, 2017). The spatial random field of the probability of presence explicitly integrates spatial autocorrelation in the probability of presence, which limits the bias in inferences due to the non-random spatial repartition of observations.

Since there were few presences for most species, our results were sensitive to the choice of prior on the variance of the spatial random effect. We tested both the inverse-gamma or a uniform distribution (the two prior forms that are available using the hSDM package; Vieilledent et al., 2014) and found that the uniform distribution provided better result. Some alternative prior choice such as the half-Cauchy are advocated in the literature (Gelman, 2006), and future work should explore if other prior choice could provide more accurate inferences. Equally, for increased confidence in predictions of rare species, several different distribution models could be compared or an ensemble model (that uses the means of several models) implemented to stabilise inferences (Araújo and New, 2007; Latimer et al., 2006; Loiselle et al., 2003).

For conservation purposes low thresholds are advised for low occurring species such as diadromous fish (Freeman and Moisen, 2008a;b; Jiménez-Valverde and Lobo, 2007). However, when lowering the threshold an increase in false positives can occur (Freeman and Moisen, 2008b). Our objective for the distribution models was to minimise false presence and absences where possible, to provide more accurate predictions whilst trying to meet conservations needs and minimise potential impacts on sea-users any spatial protection measure that might be implemented (Domisch et al., 2018; Loiselle et al., 2003; Maxwell and Jennings, 2005). Lack of understanding in species distribution can lead to inefficient spatial protection measures (Wauchope et al., 2022) and resource conflict (Probst et al., 2021).

#### 4.2. Diadromous fish river basin connectivity

The EuroDiad database is a large database that incorporates information on diadromous fish presence and abundance during their freshwater stage (Barber-O'Malley et al., 2022; Béguer et al., 2007). Using the Eurodiad database to provide

large scale understanding of river basin presence of diadromous fish, we provide a first attempt at qualitatively linking diadromous fish freshwater and marine distributions within north-eastern Atlantic waters. Although the probabilities of presence for most species were low, and gaps within the EuroDiad database evident (e.g., gaps in diadromous fish presence from Denmark and Netherlands), connectivity with their freshwater habitats appears to be visible, indicating migration pathways.

It is thought *A. alosa* no longer exists within the Greater North Sea (Baglinière et al., 2003; Wilson and Veneranta, 2019). Presences observed may have been from misidentified *A. fallax*. The high probability of presence of *A. fallax* along the east coast of the UK may be because of suitable conditions for this species as few spawning rivers have been recorded along the east coast (Aprahamian et al., 1998). For *A. agone* probabilities of presence clearly match that of their watersheds. Predicted *A. anguilla* presences closely match tagged river presences from Righton et al., (2016) despite the gaps found in the EuroDiad database. Predicted presence of *L. fluviatilis* was higher along the north-west coast of Germany and Holland. Presences have been observed here, albeit in low numbers (Admiraal et al., 1993; Pavlov et al., 2017; Thiel and Salewski, 2003). *P. marinus* predicted presence was slightly more scattered which fits with its wide distribution including into deeper waters (Elliott et al., 2021; Lança et al., 2014). The very low occurrences of *A. anguilla*, and the lamprey species is also likely to be due to inadequate sampling methods used to study these species.

Probabilities of presence of *C. ramada*, *P. flesus* and *O. eperlanus* matched relatively well their known presence from the EuroDiad database. From the EuroDiad database, there are few *S. salar* river basin presences within the North Sea. However, *S. salar* are known to occur within the North Sea and migrate Northwards to their feeding grounds (Mork et al., 2012; Rikardsen et al., 2021). Occurrences observed were most likely 1 sea winter and 2 sea winters individuals returning to their natal rivers given their size ranges. From the rivers that are monitored, *S. trutta* distribution matched well their freshwater river occupancy (ICES, 2020). The few salmonids observed may be as a result of the relatively few pelagic trawls, in addition to misreporting (ICES, 2005).

Given the coastal habitat occupancy for most of these diadromous fish and declines observed (Limburg and Waldman, 2009; Merg et al., 2020), detailed analysis of the connectivity between both habitats is essential (Flitcroft et al., 2019; Lin et al., 2017; McDowall, 2009). Unfortunately, such a model was not possible here, because it would require more detailed knowledge on their freshwater habitat occupancy (i.e., numerous outlets diadromous fish were observed within proximity to, were not contained within the EuroDiad database) and their river fidelity. Furthermore, seasonal and stage specific SDMs, which would provide more information on their ontogenetic migration movements, were not possible because of the low prevalence of the data and dispersion. Developing future research to model the connectivity between both habitats is key to improving our understanding of the pressures faced during their migrations (Flitcroft et al., 2019; Lin et al., 2017; McDowall, 2009).

#### 4.3. Gear type detectability

Even though little bycatch was observed, given a number of the diadromous fish studied here are threatened, even a small amount of bycatch may impact their populations (Dulvy et al., 2003). Furthermore, misreporting and illegal fishing is likely to remain (Elliott et al., 2020a; ICES, 2005; Stratoudakis et al., 2020; Worm et al., 2013). Bycatch detectability results from our models could help provide management advice so that gear types and areas with higher bycatch rate be avoided. Such advice can provide direct information for the MSFD and the Habitat Directive that requires the identification of gear types which may threaten protected species (1992/43/EEC; 2008/56/EC).

All species were caught by a wide range of gear types (e.g., benthic, demersal and pelagic trawls, and seine nets), despite the different gear categories being deployed at different depths. The species were probably caught by a range of gear categories because of their diel vertical migratory behaviour whilst at sea (Kristensen et al., 2018; Lança et al., 2014; Righton et al., 2016). In addition, certain gear types (e.g., trawls and static nets), can catch fish within both the demersal and pelagic water zones (He et al., 2021; Borges et al., 2008). Higher detectability from certain gear types were, however, observed. For example, shad were mainly caught by demersal mobile gear and static gear, salmonids static and pelagic mobile gear types, and *P. flesus* and *O. eperlanus* were largely captured by demersal and benthic mobile gear. These results broadly match the water column habitat occupancy these species are known to occupy and existing literature (ICES, 2005; Wilson and Veneranta, 2019). As it is thought that lampreys detach from their host upon capture (Elliott et al., 2021; Halliday, 1991), bycatch is likely to be minimal for these species.

To improve understanding of bycatch risk on a spatial level, fishing intensity by gear type should be mapped, ideally using Vessel Monitoring System data and overlaying it with the predicted distributions (Elliott et al., 2018; Quemmerais-Amice et al., 2020). Unfortunately, this information was not available to us. Additionally, since the fisheries observer data we had access to is only a small sample of the existing French fishing activity (Cornou et al., 2015), and it does not include other countries fishing effort, results would be biased.

#### 4.4. MPA relevance

To improve protection of diadromous fish at sea, numerous mechanisms could be implemented. For example, identifying areas of high bycatch risk with measures to limit bycatch within these areas, improving diadromous fish migratory pathways, and spatial protection measures (Verhelst et al., 2021; Wilson and Veneranta, 2019). Few MPAs within European waters protect marine fish and instead focus on protecting seabed features with a few exceptions (e.g., protected seabirds and marine mammals). Therefore the only protection measure that exist for vulnerable marine fish is through the removal of targeted fishing (Dureuil et al., 2018; Probst et al., 2021; Stratoudakis et al., 2016). There is, however, increasing evidence that MPAs may have positive impacts on fish populations if appropriate management measures are implemented (Davies et al., 2021; Moland et al., 2013; Probst et al., 2021; Worm et al., 2006). The value of protected areas for highly migratory species

has been questioned, but there is evidence that MPAs may have positive impacts on migratory species (Pendoley et al., 2014; Takashina and Mougi, 2014).

Our results show relatively high core area presence of protected diadromous fish within Habitat Directive MPAs. Most marine Sites of Community Importance have not been designated to protect Habitat Directive listed diadromous fish. Nonetheless, our results highlight the value of these MPAs for diadromous fish protection. Specific management measures limiting threats (e.g., limiting access to gear types with higher probability of capture) to diadromous fish within these areas, would be of benefit to their protection (Domisch et al., 2019; Stratoudakis et al., 2016). Detailed analysis of Sites of Community Importance and other categories of designated MPAs should be undertaken to evaluate their potential to protect diadromous fish.

#### 4.5. Conclusion

Understanding the spatial distribution of species and their habitats is essential for effective management and conservation (Halpern et al., 2015; Worm et al., 2009, 2005). Much research has been carried out on the conservation of freshwater diadromous fish (Drouineau et al., 2018; Merg et al., 2020; Verhelst et al., 2021). Very little research has been dedicated to the value of marine and estuary habitats for the conservation of these species and their risk of bycatch (Feunteun, 2002; Flitcroft et al., 2019; Lin et al., 2017). Here we provide an insight into the distribution and bycatch of diadromous fish at sea.

Despite diadromous fish vulnerability, targeted fishing within estuaries for numerous species remains (Aprahamian and Walker, 2008; Castelnaud, 2000; Feunteun, 2002; Kappel, 2005; Stratoudakis et al., 2020). Given the very coastal distribution of these species, more detailed analysis into such catch data (small fishing vessels targeting diadromous fish) is required. Smaller vessels and artisanal fisheries are still largely overlooked, regardless of their potential threat to diadromous fish (Beaulaton et al., 2008; Castelnaud, 2000; Stratoudakis et al., 2016). Having a better understanding of fishing pressure, combined with the outputs of the distributions and gear capture from these models, could help improve understanding of fishing impacts on diadromous fish.

Finally, it has been well acknowledged that although a large impetus for the designation of MPAs has been undertaken in the last two decades (Probst et al., 2021; Worm et al., 2006). The implementation of management measures has been slow to put in place (Dureuil et al., 2018; Probst et al., 2021; Stratoudakis et al., 2016). Furthermore, little has been undertaken to protect diadromous fish within marine ecosystems (Dureuil et al., 2018; Probst et al., 2021; Stratoudakis et al., 2016). We highlight the value of Habitat Directive MPAs that have the jurisdiction to protect diadromous fish. Limiting access of fisheries with a higher probability of capture to such areas could provide additional protection for diadromous fish and their habitats.

#### Acknowledgements

We thank the funders of the project Management of Diadromous Fish in their Environment, OFB-INRAE-Institute Agro-UPPA. We are extremely grateful to all



those who were involved in collecting and compiling the fisheries-dependent and fisheries-independent data and their funders. We are also grateful to IFREMER and the French marine fisheries and aquaculture administration (DPMA) for access to their data. Finally, thanks the reviewers for their valuable contribution.

## References

- Admiraal, W., van der Velde, G., Smit, H., Cazemier, W.G., 1993. The rivers Rhine and Meuse in The Netherlands: present state and signs of ecological recovery. *Hydrobiologia* 265, 97-128. <https://doi.org/10.1007/BF00007264>
- Alglave, B., Rivot, E., Etienne, M.-P., Woillez, M., Thorson, J.T., Vermard, Y., 2022. Combining scientific survey and commercial catch data to map fish distribution. *ICES J. Mar. Sci.* 79, 1133-1149. <https://doi.org/10.1093/icesjms/fsac032>
- Allouche, O., Tsoar, A., Kadmon, R., 2006. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). *J. Appl. Ecol.* 43, 1223-1232. <https://doi.org/10.1111/j.1365-2664.2006.01214.x>
- Almeida, P.R., 1996. Estuarine movement patterns of adult thin-lipped grey mullet, *Liza ramada* (Risso) (*Pisces*, *Mugilidae*), observed by ultrasonic tracking. *J. Exp. Mar. Bio. Ecol.* 202, 137-150. [https://doi.org/10.1016/0022-0981\(95\)00162-X](https://doi.org/10.1016/0022-0981(95)00162-X)
- Aprahamian, M., Walker, A., 2008. Status of eel fisheries, stocks and their management in England and Wales. *Knowl. Manag. Aquat. Ecosyst.* 1-7. <https://doi.org/10.1051/kmae/2009007>
- Aprahmian, M.W., Lester, S.M., Aprahamian, S.D., 1998. Shad conservation in England and Wales. R&D Technical report W110. Environment Agency: <https://www.gov.uk/government/publications/shad-conservation-in-england-and-wales> (last accessed 22/09/2022)
- Araújo, M.B., New, M., 2007. Ensemble forecasting of species distributions. *Trends Ecol. Evol.* 22, 42-47. <https://doi.org/10.1016/j.tree.2006.09.010>
- Arevalo, E., Lassalle, G., Tétard, S., Maire, A., Sauquet, E., Lambert, P., Paumier, A., Villeneuve, B., Drouineau, H., 2020. An innovative bivariate approach to detect joint temporal trends in environmental conditions: Application to large French rivers and diadromous fish. *Sci. Total Environ.* 748, 141260. <https://doi.org/10.1016/j.scitotenv.2020.141260>
- Barber-O'Malley, B., Lassalle, G., Lambert, P., Quinton, E., 2022. Dataset on European diadromous species distributions from 1750 to present time in Europe, North Africa and the Middle East. *Data Br.* 40, 107821. <https://doi.org/10.1016/j.dib.2022.107821>
- Baum, J.K., Myers, R.A., Kehler, D.G., Worm, B., Harley, S.J., Doherty, P.A., 2003. Collapse and Conservation of Shark Populations in the Northwest Atlantic. *Science*. 299, 389-392. <https://doi.org/10.1126/science.1079777>
- Bagliniere, J.L., Elie, P., 2000. Les aloses (*Alosa alosa* et *Alosa fallax* spp.) Ecobiologie et variabilité des populations. Institut National de la Recherche Agronomique, Paris. ISBN 2-7380-0906-9-ISSN :0763-1707
- Beaulaton, L., Taverny, C., Castelnaud, G., 2008. Fishing, abundance and life history traits of the anadromous sea lamprey (*Petromyzon marinus*) in Europe. *Fish. Res.* 92, 90-101. <https://doi.org/10.1016/j.fishres.2008.01.001>
- Béguier, M., Beaulaton, L., Rochard, E., 2007. Distribution and richness of diadromous fish assemblages in Western Europe: large-scale explanatory

- factors. *Ecol. Freshw. Fish* 16, 221-237. <https://doi.org/10.1111/j.1600-0633.2006.00214.x>
- Belmont, J., Miller, C., Scott, M., Wilkie, C., 2022. A new statistical approach for identifying rare species under imperfect detection. *Divers. Distrib.* 28, 882-893. <https://doi.org/10.1111/ddi.13495>
- Berg, S., Krog, C., Muus, B., Nielsen, J., Fricke, R., Berghahn, R., Neudecker, T., Wolff, W.J., 1996. Red List of lampreys and marine fishes of the Wadden Sea. *Helgoländer Meeresuntersuchungen* 50, 101-105. <https://doi.org/10.1007/BF02366178>
- Berkson, J., Thorson, J.T., 2015. The determination of data-poor catch limits in the United States: is there a better way? *ICES J. Mar. Sci.* 72, 237-242. <https://doi.org/10.1093/icesjms/fsu085>
- Bisch, A., Elliott, S.A.M., Carpentier, A., Acou, A., 2022. Modelling the distribution of vulnerable skate from fisheries dependent data using imperfect detection. *Prog. Oceanogr.* 206, 102859. <https://doi.org/10.1016/j.pocean.2022.102859>
- Bourdaud, P., Travers-Trolet, M., Vermard, Y., Cormon, X., Marchal, P., 2017. Inferring the annual, seasonal, and spatial distributions of marine species from complementary research and commercial vessels' catch rates. *ICES J. Mar. Sci.* 74, 2415-2426. <https://doi.org/10.1093/icesjms/fsx092>
- Castelnaud, G., 2000. Fishing location, fishermen and production of migratory species fisheries in the French rivers. *BFPP - Bull. Fr. la Pech. la Prot. des Milieux Aquat.* 439-458. <https://doi.org/10.1051/kmae/2001060>
- Cornou, A.S., Quinio-Scavinner, M., Delaunay, D., Dimeet, J. et al., 2015. Observations à bord des navires de pêche professionnelle. Bilan de l'échantillonnage 2014. Ifremer, Nantes
- Costa, M.J., Duarte, G., Segurado, P., Branco, P., 2021. Major threats to European freshwater fish species. *Sci. Total Environ.* 797, 149105. <https://doi.org/10.1016/j.scitotenv.2021.149105>
- Davies, B.F.R., Holmes, L., Rees, A., Attrill, M.J., Cartwright, A.Y., Sheehan, E. V., 2021. Ecosystem Approach to Fisheries Management works—How switching from mobile to static fishing gear improves populations of fished and non-fished species inside a marine-protected area. *J. Appl. Ecol.* 1-16. <https://doi.org/10.1111/1365-2664.13986>
- de Groot, R., Brander, L., van der Ploeg, S., Costanza, R., Bernard, F., Braat, L., Christie, M., Crossman, N., Ghermandi, A., Hein, L., Hussain, S., Kumar, P., McVittie, A., Portela, R., Rodriguez, L.C., ten Brink, P., van Beukering, P., 2012. Global estimates of the value of ecosystems and their services in monetary units. *Ecosyst. Serv.* 1, 50-61. <https://doi.org/10.1016/j.ecoser.2012.07.005>
- Diana, A., Matechou, E., Griffin, J.E., Buxton, A.S., Griffiths, R.A., 2021. An RShiny app for modelling environmental DNA data: accounting for false positive and false negative observation error. *Ecography (Cop.)*. 44, 1838-1844. <https://doi.org/10.1111/ecog.05718>
- Domisch, S., Friedrichs, M., Hein, T., Borgwardt, F., Wetzig, A., Jähnig, S.C., Langhans, S.D., 2018. Spatially explicit species distribution models: A missed opportunity in conservation planning? *Divers. Distrib.* 25, 758-769. <https://doi.org/10.1111/ddi.12891>
- Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber, B., Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean,

- C., Osborne, P.E., Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D., Lautenbach, S., 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography (Cop.)*. 36, 27-46. <https://doi.org/10.1111/j.1600-0587.2012.07348.x>
- Dormann, C.F., McPherson, J.M., Araújo, M.B., Bivand, R., Bolliger, J., Carl, G., Davies, R.G., Hirzel, A., Jetz, W., Daniel Kissling, W., Kühn, I., Ohlemüller, R., Peres-Neto, P.R., Reineking, B., Schröder, B., Schurr, F.M., Wilson, R., 2007. Methods to account for spatial autocorrelation in the analysis of species distributional data: a review. *Ecography (Cop.)*. 30, 609-628. <https://doi.org/10.1111/j.2007.0906-7590.05171.x>
- Drouineau, H., Carter, C., Rambonilaza, M., Beaufaron, G., Bouleau, G., Gassiat, A., Lambert, P., le Floch, S., Tétard, S., de Oliveira, E., 2018. River Continuity Restoration and Diadromous Fishes: Much More than an Ecological Issue. *Environ. Manage.* 61, 671-686. <https://doi.org/10.1007/s00267-017-0992-3>
- Dulvy, N.K., Sadovy, Y., Reynolds, J.D., 2003. Extinction vulnerability in marine populations. *Fish Fish.* 4, 25-64. <https://doi.org/10.1046/j.1467-2979.2003.00105.x>
- Dureuil, M., Boerder, K., Burnett, K.A., Froese, R., Worm, B., 2018. Elevated trawling inside protected areas undermines conservation outcomes in a global fishing hot spot. *Science*. 362, 1403-1407. <https://doi.org/10.1126/science.aau0561>
- Elliott, M., Dewailly, F., 1995. The structure and components of European estuarine fish assemblages. *Netherlands J. Aquat. Ecol.* 29, 397-417.
- Elliott, S.A.M., Bearup, D., Carpentier, A., Larivain, A., Trancart, T., Feunteun, E., 2020a. Evaluating the effectiveness of management measures on skates in a changing world. *Biol. Conserv.* 248, 108684. <https://doi.org/10.1016/j.biocon.2020.108684>
- Elliott, S.A.M., Carpentier, A., Feunteun, E., Trancart, T., 2020b. Distribution and life history trait models indicate vulnerability of skates. *Prog. Oceanogr.* 181, 102256. <https://doi.org/10.1016/j.pocean.2019.102256>
- Elliott, S.A.M., Deleys, N., Rivot, E., Acou, A., Réveillac, E., Beaulaton, L., 2021. Shedding light on the river and sea lamprey within western European marine waters. *Endanger. Species Res.* 44, 409-419. <https://doi.org/10.3354/esr01113>
- Elliott, S.A.M., Guérin, L., Pesch, R., Schmitt, P., Meakins, B., Vina-Herbon, C., González-Irusta, J.M., de la Torre, A., Serrano, A., 2018. Integrating benthic habitat indicators: Working towards an ecosystem approach. *Mar. Policy* 90, 88-94. <https://doi.org/10.1016/j.marpol.2018.01.003>
- Elliott, S.A.M., Milligan, R.J., Heath, M.R., Turrell, W.R., Bailey, D.M., 2016. Disentangling habitat concepts for demersal marine fish management. *Oceanogr. Mar. Biol. Annu. Rev.* 54, 173-191. <https://doi.org/10.1201/9781315368597>
- Engler, R., Guisan, A., Rechsteiner, L., 2004. An improved approach for predicting the distribution of rare and endangered species from occurrence and pseudo-absence data. *J. Appl. Ecol.* 41, 263-274. <https://doi.org/10.1111/j.0021-8901.2004.00881.x>
- ERSEM, 2020. NEMO-ERSEM and POLCOMS-ERSEM. Licence to use Copernicus Products. Reading, UK. [DOI.org/10.24381/cds.dcc9295c](https://doi.org/10.24381/cds.dcc9295c)

- 875 [https://cds.climate.copernicus.eu/cdsapp#!/dataset/10.24381/cds.dcc9295c?](https://cds.climate.copernicus.eu/cdsapp#!/dataset/10.24381/cds.dcc9295c?tab=overview)
- 876 [tab=overview.](https://cds.climate.copernicus.eu/cdsapp#!/dataset/10.24381/cds.dcc9295c?tab=overview)
- 877 Fauconnet, L., Trenkel, V.M., Morandeau, G., Caill-Milly, N., Rochet, M.-J., 2015.
- 878 Characterizing catches taken by different gears as a step towards evaluating
- 879 fishing pressure on fish communities. *Fish. Res.* 164, 238-248.
- 880 <https://doi.org/10.1016/j.fishres.2014.11.019>
- 881 Feunteun, E., 2002. Management and restoration of European eel population
- 882 (*Anguilla anguilla*): An impossible bargain. *Ecol. Eng.* 18, 575-591.
- 883 [https://doi.org/10.1016/S0925-8574\(02\)00021-6](https://doi.org/10.1016/S0925-8574(02)00021-6)
- 884 Fielding, A.H., Bell, J.F., 1997. A review of methods for the assessment of
- 885 prediction errors in conservation presence / absence models. *Environmental*
- 886 *Conserv.* 24, 38-49.
- 887 Flitcroft, R.L., Arismendi, I., Santelmann, M.V., 2019. A Review of Habitat
- 888 Connectivity Research for Pacific Salmon in Marine, Estuary, and Freshwater
- 889 Environments. *JAWRA J. Am. Water Resour. Assoc.* 55, 430-441.
- 890 <https://doi.org/10.1111/1752-1688.12708>
- 891 Fourcade, Y., Besnard, A.G., Secondi, J., 2018. Paintings predict the distribution of
- 892 species, or the challenge of selecting environmental predictors and evaluation
- 893 statistics. *Glob. Ecol. Biogeogr.* 27, 245-256.
- 894 <https://doi.org/10.1111/geb.12684>
- 895 Freeman, E.A., Moisen, G., 2008a. PresenceAbsence: An R Package for Presence-
- 896 Absence Model Analysis. *Journal of Statistical Software*, 23(11):1-31.
- 897 <http://www.jstatsoft.org/v23/i11>
- 898 Freeman, E.A., Moisen, G., 2008b. PresenceAbsence: An R Package for Presence-
- 899 Absence Model Analysis. *Journal of Statistical Software*, 23(11):1-31.
- 900 <http://www.jstatsoft.org/v23/i11>
- 901 Freyhof, J., Brooks, E., 2011. European Red List of freshwater fishes. Publications
- 902 Office of the European Union, Luxembourg.
- 903 [https://ec.europa.eu/environment/nature/conservation/species/redlist/dow](https://ec.europa.eu/environment/nature/conservation/species/redlist/downloads/European_freshwater_fishes.pdf)
- 904 [nloads/European\\_freshwater\\_fishes.pdf](https://ec.europa.eu/environment/nature/conservation/species/redlist/downloads/European_freshwater_fishes.pdf) (last accessed 01/07/2020).
- 905 Gelman, A., 2006. Prior distributions for variance parameters in hierarchical
- 906 models (comment on article by Browne and Draper). *Bayesian Anal.* 515-534.
- 907 <https://doi.org/10.1214/06-BA117A>
- 908 Gelman, A., Jakulin, A., Pittau, M.G., Su, Y.-S., 2008. A weakly informative default
- 909 prior distribution for logistic and other regression models. *Ann. Appl. Stat.* 2,
- 910 1360-1383. <https://doi.org/10.1214/08-AOAS191>
- 911 Guillera-Arroita, G., 2017. Modelling of species distributions, range dynamics and
- 912 communities under imperfect detection: advances, challenges and
- 913 opportunities. *Ecography (Cop.)*. 40, 281-295.
- 914 <https://doi.org/10.1111/ecog.02445>
- 915 Guisan, A., Thuiller, W., 2005. Predicting species distribution: offering more than
- 916 simple habitat models. *Ecol. Lett.* 8, 993-1009.
- 917 <https://doi.org/10.1111/j.1461-0248.2005.00792.x>
- 918 Guisan, A., Zimmermann, N.E., 2000. Predictive habitat distribution models in
- 919 ecology. *Ecol. Modell.* 135, 147-186. [https://doi.org/10.1016/S0304-](https://doi.org/10.1016/S0304-3800(00)00354-9)
- 920 [3800\(00\)00354-9](https://doi.org/10.1016/S0304-3800(00)00354-9)
- 921 Halliday, R.G., 1991. Marine distribution of the sea lamprey (*Petromyzon marinus*)
- 922 in the northwest Atlantic. *Can. J. Fish. Aquat. Sci.* 48, 832-842.
- 923 <https://doi.org/10.1139/f91-099>



- Halpern, B.S., Frazier, M., Potapenko, J., Casey, K.S., Koenig, K., Longo, C.,  
Lowndes, J.S., Rockwood, R.C., Selig, E.R., Selkoe, K.A., Walbridge, S., 2015.  
Spatial and temporal changes in cumulative human impacts on the world's  
ocean. *Nat. Commun.* 6, 1-7. <https://doi.org/10.1038/ncomms8615>
- He, P., Chopin, F., Suuronen, P., Ferro, R.S.T and Lansley, J. 2021. Classification  
and illustrated definition of fishing gears. *FAO Fisheries and Aquaculture  
Technical Paper No. 672*. Rome, FAO. <https://doi.org/10.4060/cb4966en> (last  
accessed 22/09/2020).
- Isaac, N.J.B., Jarzyna, M.A., Keil, P., Dambly, L.I., Boersch-Supan, P.H., Browning,  
E., Freeman, S.N., Golding, N., Guillera-Arroita, G., Henrys, P.A., Jarvis, S.,  
Lahoz-Monfort, J., Pagel, J., Pescott, O.L., Schmucki, R., Simmonds, E.G.,  
O'Hara, R.B., 2020. Data Integration for Large-Scale Models of Species  
Distributions. *Trends Ecol. Evol.* 35, 56-67.  
<https://doi.org/10.1016/j.tree.2019.08.006>
- ICES. 2005. Report of the Study Group on the Bycatch of Salmon in Pelagic Trawl  
Fisheries (SGBYSAL), 8-11 February 2004, Bergen, Norway. *ICES CM  
2005/ACFM:13*. 41 pp.
- ICES. 2020. Working Group with the Aim to Develop Assessment Models and  
Establish Biological Reference Points for Sea Trout (*Anadromous Salmo trutta*)  
Populations (WGTRUTTA; outputs from 2019 meeting). *ICES Scientific Reports*.  
2:59. 59 pp. <http://doi.org/10.17895/ices.pub.7431>
- Isaac, N.J.B., Jarzyna, M.A., Keil, P., Dambly, L.I., Boersch-Supan, P.H., Browning,  
E., Freeman, S.N., Golding, N., Guillera-Arroita, G., Henrys, P.A., Jarvis, S.,  
Lahoz-Monfort, J., Pagel, J., Pescott, O.L., Schmucki, R., Simmonds, E.G.,  
O'Hara, R.B., 2020. Data Integration for Large-Scale Models of Species  
Distributions. *Trends Ecol. Evol.* 35, 56-67.  
<https://doi.org/10.1016/j.tree.2019.08.006>
- Jiménez-Valverde, A., Lobo, J.M., 2007. Threshold criteria for conversion of  
probability of species presence to either-or presence-absence. *Acta  
Oecologica* 31, 361-369. <https://doi.org/10.1016/j.actao.2007.02.001>
- Kappel, C. V., 2005. Losing pieces of the puzzle: Threats to marine, estuarine, and  
diadromous species. *Front. Ecol. Environ.* 3, 275-282.  
[https://doi.org/10.1890/1540-9295\(2005\)003\[0275:LPOTPT\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2005)003[0275:LPOTPT]2.0.CO;2)
- Keith, P., Poulet, N., Denys, G., Changeux, T., Feunteun, E., et Persat, H., 2020.  
*Les poissons d'eau douce de France*. Biotope, Publications scientifiques du  
MNHN. 703 p
- Kellner, K.F., Swihart, R.K., 2014. Accounting for Imperfect Detection in Ecology: A  
Quantitative Review. *PLoS One* 9, e111436.  
<https://doi.org/10.1371/journal.pone.0111436>
- Korpinen, S., Laamanen, L., Bergström, L., Nurmi, M., Andersen, J.H., Haapaniemi,  
J., Harvey, E.T., Murray, C.J., Peterlin, M., Kallenbach, E., Klančnik, K.,  
Stein, U., Tunesi, L., Vaughan, D., Reker, J., 2021. Combined effects of  
human pressures on Europe's marine ecosystems. *Ambio* 50, 1325-1336.  
<https://doi.org/10.1007/s13280-020-01482-x>
- Kristensen, M., Righton, D., del Villar-Guerra, D., Baktoft, H., Aarestrup, K., 2018.  
Temperature and depth preferences of adult sea trout *Salmo trutta* during  
the marine migration phase. *Mar. Ecol. Prog. Ser.* 599, 209-224.  
<https://doi.org/10.3354/meps12618>

- 972 Lambert, C., Authier, M., Dorémus, G., Laran, S., Panigada, S., Spitz, J., Van  
973 Canneyt, O., Ridoux, V., 2020. Setting the scene for Mediterranean litterscape  
974 management: The first basin-scale quantification and mapping of floating  
975 marine debris. *Environ. Pollut.* 263, 114430.  
976 <https://doi.org/10.1016/j.envpol.2020.114430>
- 977 Lança, M.J., Machado, M., Mateus, C.S., Lourenço, M., Ferreira, A.F., Quintella,  
978 B.R., Almeida, P.R., 2014. Investigating Population Structure of Sea Lamprey  
979 (*Petromyzon marinus*, L.) in Western Iberian Peninsula Using Morphological  
980 Characters and Heart Fatty Acid Signature Analyses. *PLoS One* 9, e108110.  
981 <https://doi.org/10.1371/journal.pone.0108110>
- 982 Lassalle, G., Béguer, M., Beaulaton, L., Rochard, E., 2008. Diadromous fish  
983 conservation plans need to consider global warming issues: An approach using  
984 biogeographical models. *Biol. Conserv.* 141, 1105-1118.  
985 <https://doi.org/10.1016/j.biocon.2008.02.010>
- 986 Lassalle, G., 2021. EuroDiad version 4.0, <https://doi.org/10.15454/IVVAIC>, Portail  
987 Data INRAE, V2, UNF:6:X15FgpfOWssoJvGSg0wyoA== [fileUNF]
- 988 Latimer, A.M., Wu, S., Gelfand, A.E., Silander Jr., J.A., 2006. Building Statistical  
989 Models To Analyze Species Distributions. *Ecol. Appl.* 16, 33-50.  
990 <https://doi.org/10.1890/04-0609>
- 991 Leathwick, J.R., Rowe, D., Richardson, J., Elith, J., Hastie, T., 2005. Using  
992 multivariate adaptive regression splines to predict the distributions of New  
993 Zealand's freshwater diadromous fish. *Freshw. Biol.* 50, 2034-2052.  
994 <https://doi.org/10.1111/j.1365-2427.2005.01448.x>
- 995 Legrand, M., Briand, C., Buisson, L., Artur, G., Azam, D., Baisez, A., Barracou, D.,  
996 Bourré, N., Carry, L., Caudal, A.-L., Charrier, F., Corre, J., Croguennec, E.,  
997 Der Mikaélian, S., Josset, Q., Le Gurun, L., Schaeffer, F., Laffaille, P., 2020.  
998 Contrasting trends between species and catchments in diadromous fish counts  
999 over the last 30 years in France. *Knowl. Manag. Aquat. Ecosyst.* 2020-Janua,  
1000 7. <https://doi.org/10.1051/kmae/2019046>
- 1001 Leroy, B., Delsol, R., Hugueny, B., Meynard, C.N., Barhoumi, C., Barbet-Massin, M.,  
1002 Bellard, C., 2018. Without quality presence-absence data, discrimination  
1003 metrics such as TSS can be misleading measures of model performance. *J.*  
1004 *Biogeogr.* 45, 1994-2002. <https://doi.org/10.1111/jbi.13402>
- 1005 Limburg, K.E., Waldman, J.R., 2009. Dramatic Declines in North Atlantic  
1006 Diadromous Fishes. *Bioscience* 59, 955-965.  
1007 <https://doi.org/10.1525/bio.2009.59.11.7>
- 1008 Lin, H., Bush, A., Linke, S., Possingham, H.P., Brown, C.J., 2017. Climate change  
1009 decouples marine and freshwater habitats of a threatened migratory fish.  
1010 *Divers. Distrib.* 23, 751-760. <https://doi.org/10.1111/ddi.12570>
- 1011 Loiselle, B.A., Howell, C.A., Graham, C.H., Goerck, J.M., Brooks, T., Smith, K.G.,  
1012 Williams, P.H., 2003. Avoiding Pitfalls of Using Species Distribution Models in  
1013 Conservation Planning. *Conserv. Biol.* 17, 1591-1600.  
1014 <https://doi.org/10.1111/j.1523-1739.2003.00233.x>
- 1015 Lomba, A., Pellissier, L., Randin, C., Vicente, J., Moreira, F., Honrado, J., Guisan,  
1016 A., 2010. Overcoming the rare species modelling paradox: A novel hierarchical  
1017 framework applied to an Iberian endemic plant. *Biol. Conserv.* 143, 2647-  
1018 2657. <https://doi.org/10.1016/j.biocon.2010.07.007>
- 1019 Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H., Cooke, R.G., Kay, M.C.,  
1020 Kidwell, S.M., Kirby, M.X., Peterson, C.H., Jackson, J.B.C., 2006. Depletion,

- 1021 Degradation, and Recovery Potential of Estuaries and Coastal Seas. *Science*.  
1022 312, 1806-1809. <https://doi.org/10.1126/science.1128035>
- 1023 MacKenzie, D.I., Nichols, J.D., Lachman, G.B., Droege, S., Royle, A.A., Langtimm,  
1024 C.A., 2002. Estimating site occupancy rates when detection probabilities are  
1025 less than one. *Ecology* 83, 2248-2255. [https://doi.org/10.1890/0012-](https://doi.org/10.1890/0012-9658(2002)083[2248:ESORWD]2.0.CO;2)  
1026 9658(2002)083[2248:ESORWD]2.0.CO;2
- 1027 Maes, J., Stevens, M., Breine, J., 2007. Modelling the migration opportunities of  
1028 diadromous fish species along a gradient of dissolved oxygen concentration in  
1029 a European tidal watershed. *Estuar. Coast. Shelf Sci.* 75, 151-162.  
1030 <https://doi.org/10.1016/j.ecss.2007.03.036>
- 1031 Maxwell, D., Jennings, S., 2005. Power of monitoring programmes to detect decline  
1032 and recovery of rare and vulnerable fish. *J. Appl. Ecol.* 42, 25-37.  
1033 <https://doi.org/10.1111/j.1365-2664.2005.01000.x>
- 1034 McDowall, R.M., 2009. Making the Best of Two Worlds : Diadromy in the Evolution,  
1035 Ecology, and Conservation of Aquatic Organisms. *Am. Fish. Soc. Symp.* 69, 1-  
1036 22.
- 1037 Merg, M.-L., Dézerald, O., Kreutzenberger, K., Demski, S., Reyjol, Y., Usseglio-  
1038 Polatera, P., Belliard, J., 2020. Modeling diadromous fish loss from historical  
1039 data: Identification of anthropogenic drivers and testing of mitigation  
1040 scenarios. *PLoS One* 15, e0236575.  
1041 <https://doi.org/10.1371/journal.pone.0236575>
- 1042 Meyer, C.F.J., Aguiar, L.M.S., Aguirre, L.F., Baumgarten, J., Clarke, F.M., Cosson,  
1043 J.F., Villegas, S.E., Fahr, J., Faria, D., Furey, N., Henry, M., Hodgkison, R.,  
1044 Jenkins, R.K.B., Jung, K.G., Kingston, T., Kunz, T.H., Cristina Macswiney  
1045 Gonzalez, M., Moya, I., Patterson, B.D., Pons, J.M., Racey, P.A., Rex, K.,  
1046 Sampaio, E.M., Solari, S., Stoner, K.E., Voigt, C.C., von Staden, D., Weise,  
1047 C.D., Kalko, E.K.V., 2011. Accounting for detectability improves estimates of  
1048 species richness in tropical bat surveys. *J. Appl. Ecol.* 48, 777-787.  
1049 <https://doi.org/10.1111/j.1365-2664.2011.01976.x>
- 1050 Moland, E., Olsen, E.M., Knutsen, H., Garrigou, P., Espeland, S.H., Kleiven, A.R.,  
1051 Andre, C., Knutsen, J.A., 2013. Lobster and cod benefit from small-scale  
1052 northern marine protected areas: inference from an empirical before-after  
1053 control-impact study. *Proc. R. Soc. B Biol. Sci.* 280, 1-10.  
1054 <https://doi.org/10.1098/rspb.2012.2679>
- 1055 Moriarty, M., Sethi, S.A., Pedreschi, D., Smeltz, T.S., McGonigle, C., Harris, B.P.,  
1056 Wolf, N., Greenstreet, S.P.R., 2020. Combining fisheries surveys to inform  
1057 marine species distribution modelling. *ICES J. Mar. Sci.* 77, 539-552.  
1058 <https://doi.org/10.1093/icesjms/fsz254>
- 1059 Nieto, A., Ralph, G.M., Comeros-Raynal, M.T., Kemp, J., Garcia Criado, M., Allen,  
1060 D.J., Dulvy, N.K., Walls, R.H.L., Russell, B., Pollard, D., et al., 2015.  
1061 European Red List of marine fishes. Luxembourg: Publications Office of the  
1062 European Union.  
1063 [https://ec.europa.eu/environment/nature/conservation/species/redlist/dow](https://ec.europa.eu/environment/nature/conservation/species/redlist/downloads/European_marine_fishes.pdf)  
1064 nloads/European\_marine\_fishes.pdf (last accessed 01/07/2020)
- 1065 Mork, K.A., Gilbey, J., Hansen, L.P., Jensen, A.J., Jacobsen, J.A., Holm, M., Holst,  
1066 J.C., Ó Maoiléidigh, N., Vikebø, F., McGinnity, P., Melle, W., Thomas, K.,  
1067 Verspoor, E., Wennevik, V., 2012. Modelling the migration of post-smolt  
1068 Atlantic salmon (*Salmo salar*) in the Northeast Atlantic. *ICES J. Mar. Sci.* 69,  
1069 1616-1624. <https://doi.org/10.1093/icesjms/fss108>

- Northrup, J.M., Gerber, B.D., 2018. A comment on priors for Bayesian occupancy models. PLoS One 13, e0192819.  
<https://doi.org/10.1371/journal.pone.0192819>
- Pavlov, D.S., Zvezdin, A.O., Kostin, V. V., Tsimbalov, I.A., Kucheryavyy, A. V., 2017. Temporal characteristics of downstream migration of smolts of the European river lamprey *Lampetra fluviatilis* in the Chernaya River. Biol. Bull. 44, 290-295. <https://doi.org/10.1134/S1062359017030062>
- Pendoley, K.L., Schofield, G., Whittock, P.A., Ierodiaconou, D., Hays, G.C., 2014. Protected species use of a coastal marine migratory corridor connecting marine protected areas. Mar. Biol. 161, 1455-1466.  
<https://doi.org/10.1007/s00227-014-2433-7>
- Prince, J., Hordyk, A., 2019. What to do when you have almost nothing: A simple quantitative prescription for managing extremely data-poor fisheries. Fish Fish. 20, 224-238. <https://doi.org/10.1111/faf.12335>
- Probst, W.N., Stelzenmüller, V., Rambo, H., Moriarty, M., Greenstreet, S.P.R., 2021. Identifying core areas for mobile species in space and time: A case study of the demersal fish community in the North Sea. Biol. Conserv. 254, 108946. <https://doi.org/10.1016/j.biocon.2020.108946>
- Quemmerais-Amice, F., Barrere, J., La Rivière, M., Contin, G., Bailly, D., 2020. A Methodology and Tool for Mapping the Risk of Cumulative Effects on Benthic Habitats. Front. Mar. Sci. 7. <https://doi.org/10.3389/fmars.2020.569205>
- Righton, D., Westerberg, H., Feunteun, E., Økland, F., Gargan, P., Amilhat, E., Metcalfe, J., Lobon-Cervia, J., Sjöberg, N., Simon, J., Acou, A., Vedor, M., Walker, A., Trancart, T., Brämick, U., Aarestrup, K., 2016. Empirical observations of the spawning migration of European eels: The long and dangerous road to the Sargasso Sea. Sci. Adv. 2, e1501694.  
<https://doi.org/10.1126/sciadv.1501694>
- Rikardsen, A.H., Righton, D., Strøm, J.F., Thorstad, E.B., Gargan, P., Sheehan, T., Økland, F., Chittenden, C.M., Hedger, R.D., Næsje, T.F., Renkawitz, M., Sturlaugsson, J., Caballero, P., Baktoft, H., Davidsen, J.G., Halttunen, E., Wright, S., Finstad, B., Aarestrup, K., 2021. Redefining the oceanic distribution of Atlantic salmon. Sci. Rep. 11, 12266.  
<https://doi.org/10.1038/s41598-021-91137-y>
- Royle, J.A., Link, W.A., 2006. Generalized site occupancy models allowing for false positive and false negative errors. Ecology 87, 835-841.  
[https://doi.org/10.1890/0012-9658\(2006\)87\[835:GSOMAF\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[835:GSOMAF]2.0.CO;2)
- Simmonds, E.G., Jarvis, S.G., Henrys, P.A., Isaac, N.J.B., O'Hara, R.B., 2020. Is more data always better? A simulation study of benefits and limitations of integrated distribution models. Ecography (Cop.). 43, 1413-1422.  
<https://doi.org/10.1111/ecog.05146>
- Spiegelhalter, D.J., Best, N.G., Carlin, B.P., van der Linde, A., 2002. Bayesian measures of model complexity and fit. J. R. Stat. Soc. Ser. B (Statistical Methodol. 64, 583-639. <https://doi.org/10.1111/1467-9868.00353>
- Stratoudakis, Y., Correia, C., Belo, A.F., de Almeida, P.R., 2020. Improving participated management under poor fishers' organization: Anadromous fishing in the estuary of Mondego River, Portugal. Mar. Policy 119, 104049.  
<https://doi.org/10.1016/j.marpol.2020.104049>
- Stratoudakis, Y., Mateus, C.S., Quintella, B.R., Antunes, C., Raposo de Almeida, P., 2016. Exploited anadromous fish in Portugal: Suggested direction for



- conservation and management. Mar. Policy 73, 92-99.  
<https://doi.org/10.1016/j.marpol.2016.07.031>
- Takashina, N., Mougi, A., 2014. Effects of marine protected areas on overfished fishing stocks with multiple stable states. J. Theor. Biol. 341, 64-70.  
<https://doi.org/10.1016/j.jtbi.2013.09.027>
- Taverny, C., Elie, P., 2001. Répartition spatio-temporelle de la grande alose *Alosa alosa* (Linne, 1766) et de l'Alose feinte *Alosa fallax* (Lacépède, 1803) dans le Golfe de Gascogne. Bull. Français la Pêche la Piscic. 803-821.  
<https://doi.org/10.1051/kmae:2001020>
- Taverny, C., Lassalle, G., Ortusi, I., Roqueplo, C., Lepage, M., Lambert, P., 2012. From shallow to deep waters: habitats used by larval lampreys (genus *Petromyzon* and *Lampetra*) over a western European basin. Ecol. Freshw. Fish 21, 87-99. <https://doi.org/10.1111/j.1600-0633.2011.00526.x>
- Teruzzi, A., Di Biagio, V., Feudale, L., Bolzon, G., Lazzari, P., Salon, S., Di Biagio, V., Coidessa, G., and Cossarini, G., 2021. Mediterranean Sea Biogeochemical Reanalysis (CMEMS MED-Biogeochemistry, MedBFM3 system) (Version 1) [Data set]. Copernicus Monitoring Environment Marine Service (CMEMS).  
[https://doi.org/10.25423/CMCC/MEDSEA\\_MULTIYEAR\\_BGC\\_006\\_008\\_MEDBFM3](https://doi.org/10.25423/CMCC/MEDSEA_MULTIYEAR_BGC_006_008_MEDBFM3)
- Thiel, R., Salewski, V., 2003. Verteilung und Wanderung von Neunaugen im Elbeästuar (Deutschland). Limnologica 33, 214-226.  
[https://doi.org/10.1016/S0075-9511\(03\)80015-4](https://doi.org/10.1016/S0075-9511(03)80015-4)
- Trancart, T., Rochette, S., Acou, A., Lasne, E., Feunteun, E., 2014. Modeling marine shad distribution using data from French bycatch fishery surveys. Mar. Ecol. Prog. Ser. 511, 181-192. <https://doi.org/10.3354/meps10907>
- Verhelst, P., Reubens, J., Buysse, D., Goethals, P., Van Wichelen, J., Moens, T., 2021. Toward a roadmap for diadromous fish conservation: the Big Five considerations. Front. Ecol. Environ. fee.2361.  
<https://doi.org/10.1002/fee.2361>
- Vieilledent, G., Merow, C., Guelat, J., Latimer, A.M., Kery, M., Gelfand, A.E., Wilson, A.M., Mortier, F., Silander Jr., J.A., 2014. hSDM: Hierarchical Bayesian Species Distribution Models. R package version 1.4. <https://CRAN.R-project.org/package=hSDM>
- Waldman, J.R., Quinn, T.P., 2022. North American diadromous fishes: Drivers of decline and potential for recovery in the Anthropocene. Sci. Adv. 8.  
<https://doi.org/10.1126/sciadv.abl5486>
- Wauchope, H.S., Jones, J.P.G., Geldmann, J., Simmons, B.I., Amano, T., Blanco, D.E., Fuller, R.A., Johnston, A., Langendoen, T., Mundkur, T., Nagy, S., Sutherland, W.J., 2022. Protected areas have a mixed impact on waterbirds, but management helps. Nature 605, 103-107.  
<https://doi.org/10.1038/s41586-022-04617-0>
- Wilson, K., and Veneranta, L. (Eds). 2019. Data-limited diadromous species - review of European status. ICES Cooperative Research Report No. 348. 273 pp.  
<http://doi.org/10.17895/ices.pub.5253>
- Worm, B., Barbier, E.B., Beaumont, N., Duffy, J.E., Folke, C., Halpern, B.S., Jackson, J.B.C., Lotze, H.K., Micheli, F., Palumbi, S.R., Sala, E., Selkoe, K.A., Stachowicz, J.J., Watson, R., 2006. Impacts of Biodiversity Loss on Ocean Ecosystem Services. Science (80-. ). 314, 787-790.  
<https://doi.org/10.1126/science.1132294>

1167 Worm, B., Davis, B., Ketteimer, L., Ward-Paige, C.A., Chapman, D., Heithaus, M.R.,  
 1168 Kessel, S.T., Gruber, S.H., 2013. Global catches, exploitation rates, and  
 1169 rebuilding options for sharks. *Mar. Policy* 40, 194-204.  
 1170 <https://doi.org/10.1016/j.marpol.2012.12.034>  
 1171 Worm, B., Hilborn, R., Baum, J.K., Branch, T.A., Collie, J.S., Costello, C., Fogarty,  
 1172 M.J., Fulton, E.A., Hutchings, J.A., Jennings, S., Jensen, O.P., Lotze, H.K.,  
 1173 Mace, P.M., McClanahan, T.R., Minto, C., Palumbi, S.R., Parma, A.M., Ricard,  
 1174 D., Rosenberg, A.A., Watson, R., Zeller, D., 2009. Rebuilding global fisheries.  
 1175 *Science*. 325, 578-85. <https://doi.org/10.1126/science.1173146>  
 1176 Worm, B., Sadow, M., Oschlies, A., Lotze, H., Myers, R., 2005. Global patterns of  
 1177 predator diversity in the open oceans. *Science*. 309, 1365-9.  
 1178 <https://doi.org/10.1126/science.1113399>

